À des fins de recherche uniquement

Anticorps Monoclonal anti-CD226

Numéro de catalogue: CR-65247



Informations de base

Numéro de catalogue:

100 tests , 5 μ l/test

CR-65247

Taille:

Hôte:

IgG1, kappa

Numéro d'acquisition GenBank:

BC074787

Méthode de purification:

Identification du gène (NCBI):

CloneNo.:

10666

11A8

Excitation/Emission maxima

Mouse Isotype:

Nom complet: CD226 molecule MW calculé 336 aa, 39 kDa

wavelengths: 592 nm / 611 nm

Applications

Applications testées:

FC

Spécificité de l'espèce:

Humain

Informations générales

CD226 (DNAM-1) is a ~65 kDa glycoprotein expressed on the surface of NK cells, platelets, monocytes and a subset of T cells. It is a member of the Ig-superfamily containing 2 Ig-like domains of the V-set. CD226 mediates cellular adhesion of platelets and megakaryocytic cells to vascular endothelial cells. The protein also plays a role in megakaryocytic cell maturation. Interactions of CD226 and its ligands, CD155 and CD112, induce NK and T cellmediated cytotoxicity and cytokine secretion (PMID: 15039383).


Stockage

Stockage:


Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 0.09% sodium azide and 0.5% BSA.

Données de validation sélectionnées



1X10^6 human PBMCs were surface co-stained with Atlantic Blue Anti-Human CD3 and 5 ul Cardinal Red™ Anti-Human CD226 (CR-65247, Clone:11A8) or Mouse IgG1 Isotype Control. Cells were not fixed. Lymphocytes were gated.



1X10^6 human PBMCs were surface stained with 5 ul Cardinal Red™ Anti-Human CD226 (CR-65247, Clone:11A8) or Mouse IgG1 Isotype Control. Cells were not fixed. Lymphocytes were gated.