À des fins de recherche uniquement

Anticorps Monoclonal anti-CD64

Numéro de catalogue: CR-65253



Informations de base

Numéro de catalogue:

Numéro d'acquisition GenBank:

BC032634

Méthode de purification: Purification par affinité

CR-65253 Taille:

Identification du gène (NCBI): 2209

CloneNo.:

100 tests , 5 μ l/test

Nom complet:

Excitation/Emission maxima

Hôte: Mouse

Fc fragment of IgG, high affinity Ia, receptor (CD64)

wavelengths: 592 nm / 611 nm

Isotype: IgG1, kappa

MW calculé 374 aa, 43 kDa

Applications

Applications testées:

Spécificité de l'espèce:

Humain

Informations générales

Fcy receptor comprise a multigene family of integral membrane glycoproteins that exhibit complex activation or $inhibitory\ effects\ on\ cell\ functions\ after\ aggregation\ by\ complexed\ immunoglobulin\ G\ (IgG)\ (PMID:\ 17005690\).$ CD64, also known as Fcy RIA, is a high-affinity receptor for the Fc region of IgG. It is expressed by monocytes/macrophages, activated neutrophils, dendritic cells, and early myeloid cells (PMID: 23293080; 19642859; 7680917). CD64 functions in both innate and adaptive immune responses.


Stockage

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.


Tampon de stockage:

PBS with 0.09% sodium azide and 0.5% BSA.

Données de validation sélectionnées



1X10^6 human PBMCs were surface stained with 5 ul Cardinal Red™ Anti-Human CD64 (CR-65253, Clone:10.1) (red) or unstained. Cells were not fixed. Monocytes were gated.



1X10^6 human PBMCs were surface stained with 5 ul Cardinal Red™ Anti-Human CD64 (CR-65253, Clone:10.1) or unstained. Cells were not fixed. Monocytes were gated.