

À des fins de recherche uniquement

# Anticorps Monoclonal anti-CD55

Numéro de catalogue: **CR-65271**



## Informations de base

<b>Numéro de catalogue:</b> CR-65271	<b>Numéro d'acquisition GenBank:</b> BC001288	<b>Méthode de purification:</b> Purification par affinité
<b>Taille:</b> 100tests , 5 ul/test	<b>Identification du gène (NCBI):</b> 1604	<b>CloneNo.:</b> JS11
<b>Hôte:</b> Mouse	<b>Nom complet:</b> CD55 molecule, decay accelerating factor for complement (Cromer blood group)	<b>Excitation/Emission maxima wavelengths:</b> 592 nm / 611 nm
<b>Isotype:</b> IgG1	<b>MW calculé</b> 41 kDa	

## Applications

**Applications testées:**  
FC

**Spécificité de l'espèce:**  
Humain

## Informations générales

CD55, also known as DAF, is a glycosylphosphatidylinositol (GPI)-anchored surface glycoprotein that is widely distributed on blood, stroma, epithelial, and endothelial cells (PMID: 7517044; 29503741). It can also exist as a soluble form in plasma, urine, saliva, tears, and synovial fluids (PMID: 29503741). CD55 is a complement regulatory protein (PMID: 2469439; 7517044). It inhibits formation of the C3 convertases through binding to C3b and C4b. It also binds the alternate pathway convertase C3bBb, the classical pathway convertase and C4b2a to accelerate their decay (PMID: 17289551). CD55 also serves as a receptor for coxsackieviruses B1, B3, and B5 and several enteroviruses (PMID: 7538177; 7517044).

## Stockage

**Stockage:**  
Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

**Tampon de stockage:**  
PBS with 0.09% sodium azide and 0.5% BSA.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**

## Données de validation sélectionnées



1X10<sup>6</sup> human PBMCs were surface stained with 5 ul Cardinal Red™ Anti-Human CD55 (CR-65271, Clone: JS11) (red) or Mouse IgG1 Isotype Control. Cells were not fixed. Lymphocytes were gated.



1X10<sup>6</sup> human PBMCs were surface stained with 5 ul Cardinal Red™ Anti-Human CD55 (CR-65271, Clone: JS11) or Mouse IgG1 Isotype Control. Cells were not fixed. Lymphocytes were gated.