

À des fins de recherche uniquement

Anticorps Monoclonal anti-IL-12/IL-23 p40



Numéro de catalogue: FITC-65097

Informations de base

Numéro de catalogue:

FITC-65097

Taille:

100ug, 500 µg/ml

Hôte:

Rat

Isotype:

IgG2a, kappa

Numéro d'acquisition GenBank:

BC103608

Identification du gène (NCBI):

16160

Nom complet:

interleukin 12b

Méthode de purification:

Purification par affinité

CloneNo.:

C17.8

Excitation/Emission maxima wavelengths:

495 nm / 524 nm

Applications

Applications testées:

FC (Intra)

Spécificité de l'espèce:

souris

Informations générales

Interleukin (IL)-12 and IL-23 are heterodimeric cytokines that share a common p40 subunit (PMID: 11114383). IL-12 is composed of the IL-12 p40 subunit linked to the IL-12 p35 subunit, and the heterodimer signals through the IL-12 receptor (IL-12R), which comprises the IL-12R β 1 and IL-12R β 2 subunits. IL-23 is composed of the IL-23 p19 subunit and the IL-12 p40 (IL-12/23p40) subunit, which signals through IL-23R and IL-12R β 1 (PMID: 11114383; 26121196). IL-12/IL-23 p40 also exists as a monomer and as a homodimer which can act as a potent IL-12 antagonist (PMID: 8958912; 18783467). IL-12/IL-23 p40 is produced by antigen-presenting cells, such as dendritic cells (DCs), monocytes, macrophages, neutrophils and, to a lesser extent, B cells (PMID: 20476918).

Stockage

Stockage:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

Tampon de stockage:

PBS with 0.09% sodium azide.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



1X10⁶ IFN- γ primed Balb/c mouse peritoneal macrophages were unstimulated or stimulated with LPS in the presence of a protein transport inhibitor and then intracellularly stained with 0.5 ug FITC Plus Anti-Mouse IL-12/IL-23 p40 (FITC-65097, Clone:C17.8). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).