

À des fins de recherche uniquement

Anticorps Monoclonal anti-CD8

Numéro de catalogue: FITC-65146



Informations de base

Numéro de catalogue:

FITC-65146

Taille:

100 tests, 5 µl/test

Hôte:

Mouse

Isotype:

IgG1, kappa

Numéro d'acquisition GenBank:

BC025715

Identification du gène (NCBI):

925

Nom complet:

CD8a molecule

MW calculé

235 aa, 26 kDa

Méthode de purification:

Purification par affinité

CloneNo.:

SK1

Excitation/Emission maxima
wavelengths:

494 nm / 520 nm

Applications

Applications testées:

FC

Spécificité de l'espèce:

Humain

Informations générales

CD8 is a transmembrane glycoprotein composed of two disulfide-linked chains. It can be present as a homodimer of CD8 α or as a heterodimer of CD8 α and CD8 β (PMID: 3264320; 8253791). CD8 is found on most thymocytes. The majority of class I-restricted T cells express mostly the CD8 $\alpha\beta$ heterodimer while CD8 $\alpha\alpha$ homodimers alone have been found on some gut intraepithelial T cells, on some T cell receptor (TCR) $\gamma\delta$ T cells and on NK cells (PMID: 2111591; 1831127; 8420975). CD8 acts as a co-receptor that binds to MHC class-I and participates in cytotoxic T cell activation (PMID: 8499079). During T cell development, CD8 is required for positive selection of CD4-/CD8+ T cells (PMID: 1968084).

Stockage

Stockage:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

Tampon de stockage:

Tampon à base de phosphate avec azoture de sodium à 0,09 % et gélatine à 0,1 %, pH 7,2.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



1X10⁶ human peripheral blood lymphocytes were surface stained with 5.00 ul/test APC-anti-human CD3 (APC-65151, Clone: UCHT1) and 5.00 ul/test FITC-anti-human CD8 (FITC-65146, Clone: SK1). Cells were not fixed.



1X10⁶ human peripheral blood lymphocytes were surface stained with 5.00 ul/test APC-anti-human CD3 (APC-65151, Clone: UCHT1) and 0.125 ug FITC-mouse IgG1 isotype control. Cells were not fixed.