

À des fins de recherche uniquement

# Anticorps Monoclonal anti-CD279

Numéro de catalogue:**FITC-65168**



## Informations de base

Numéro de catalogue:	FITC-65168	Numéro d'acquisition GenBank:	BC074740	Méthode de purification:
Taille:	100 tests, 4 µl/test	Identification du gène (NCBI):	5133	L'anticorps purifié est conjugué à l'isothiocyanate de fluorescéine (FITC) dans des conditions optimales.
Hôte:	Mouse	Nom complet:	programmed cell death 1	Le conjugué est purifié par chromatographie d'exclusion de taille.
Isotype:	IgG1, kappa	MW calculé	288 aa, 32 kDa	CloneNo.:
				EH12.2H7
				Excitation/Emission maxima wavelengths:
				498 nm / 526 nm

## Applications

### Applications testées:

FC

### Spécificité de l'espèce:

Humain, primates non humains

## Informations générales

Programmed cell death 1 (PD-1, also known as CD279) is an immunoinhibitory receptor that belongs to the CD28/CTLA-4 subfamily of the Ig superfamily. It is a 288 amino acid (aa) type I transmembrane protein composed of one Ig superfamily domain, a stalk, a transmembrane domain, and an intracellular domain containing an immunoreceptor tyrosine-based inhibitory motif (ITIM) as well as an immunoreceptor tyrosine-based switch motif (ITSM) (PMID: 18173375). PD-1 is expressed during thymic development and is induced in a variety of hematopoietic cells in the periphery by antigen receptor signaling and cytokines (PMID: 20636820). Engagement of PD-1 by its ligands PD-L1 or PD-L2 transduces a signal that inhibits T-cell proliferation, cytokine production, and cytolytic function (PMID: 19426218). It is critical for the regulation of T cell function during immunity and tolerance. Blockade of PD-1 can overcome immune resistance and also has been shown to have antitumor activity (PMID: 22658127; 23169436).

## Stockage

### Stockage:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

### Tampon de stockage:


PBS avec azoture de sodium à 0,1 %.

For technical support and original validation data for this product please contact:  
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free  
in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com  
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Données de validation sélectionnées



1X10<sup>6</sup> human peripheral blood mononuclear cells (PBMCs) were surface stained with APC Anti-Human CD3 (APC-65151, clone UCHT1) and 4.00  $\mu$ l/test FITC Anti-Human CD279 (FITC-65168, Clone:EH12.2H7). Cells were not fixed.



1X10<sup>6</sup> human peripheral blood mononuclear cells (PBMCs) were surface stained with APC Anti-Human CD3 (APC-65151, clone UCHT1) and FITC mouse IgG1 isotype control antibody. Cells were not fixed.