À des fins de recherche uniquement

## Anticorps Monoclonal anti-CD28

Numéro de catalogue:PE-65099





Informations de base	Numéro de catalogue: PE-65099	Numéro d'acquisition GenBank: BC093698	Méthode de purification: Purification par affinité
	Taille: 100tests , 5 µl/test	Identification du gène (NCBI): 940	CloneNo.: CD28.2
	Hôte: Mouse Isotype: IgG1, kappa	Nom complet: CD28 molecule MW calculé 220 aa, 25 kDa	Excitation/Emission maxima wavelengths: 496 nm, 565 nm / 578 nm
Spécificité de l'espèce: Humain			
Informations générales	CD28 (T-cell-specific surface glycoprotein CD28), also known as T44 and Tp44, is a 44 kD disulfide-linked homodimeric type I glycoprotein (PMID: 2162180). It is a member of the immunoglobulin superfamily and is expressed on most T lineage cells, NK cell subsets, and plasma cells (PMID: 2162180, 8386518). CD28 may affect in vivo immune responses by functioning both as a cell adhesion molecule linking B and T lymphocytes and as the surface component of a novel signal transduction pathway (PMID: 2162180, 3021470). CD28 binds both CD80 and CD86 with a highly conserved motif MYPPY in the CDR3-like loop (PMID: 15696168, 7964482). CD28 is considered a major co-stimulatory molecule, inducing T lymphocyte activation and IL-2 synthesis, and preventing cell death (PMID: 1348520).		
Stockage	Stockage: Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment. Tampon de stockage: PBS with 0.09% sodium azide.		

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free<br/>in USA), or 1(312) 455-8498 (outside USA)E: proteintech@ptglab.comW: ptglab.comW: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Données de validation sélectionnées





1X10<sup>^6</sup> human PBMCs were surface co-stained with CD3 and 5 ul PE Anti-Human CD28 (PE-65099, Clone:CD28.2) or Mouse IgG2a Isotype Control. Cells were not fixed. Lymphocytes were gated.

1X10<sup>^6</sup> human PBMCs were surface stained with 5 ul PE Anti-Human CD28 (PE-65099, Clone:CD28.2) or Mouse IgG2a Isotype Control. Cells were not fixed. Lymphocytes were gated.