## À des fins de recherche uniquement

## Anticorps Monoclonal anti-CD19

Numéro de catalogue:PE-65110



Informations de base

Numéro de catalogue:

PE-65110

Taille:

100 tests , 5  $\mu$ l/test

Hôte: Mouse Isotype: IgG1, kappa Numéro d'acquisition GenBank:

BC006338 Identification du gène (NCBI):

Nom complet: CD19 molecule MW calculé

556 aa, 61 kDa

Méthode de purification: Purification par affinité

CloneNo.: HIB19

Excitation/Emission maxima wavelengths:

496 nm, 565 nm / 578 nm

**Applications** 

Applications testées:

FC

Spécificité de l'espèce:

Humain

Informations générales

CD19 is a 95 kDa type I transmembrane glycoprotein belonging to the immunoglobulin superfamily (PMID: 2472450). It is expressed by B cells and follicular dendritic cells. CD19 is up-regulated at the step of B-lineage commitment during the differentiation of the hematopoietic stem cell, it remains on during subsequent stages of differentiation until finally down-regulated during terminal differentiation into plasma cells (PMID: 8528044). CD19 is involved in B cell development, activation and differentiation. It is the dominant component for the signaling complex on B cells that includes CD21 (CR2), CD81 (TAPA-1) and CD225 and acts as a critical co-receptor for BCR signal transduction (PMID: 23210908).

Stockage


Stockage:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.


Tampon de stockage:

PBS with 0.09% sodium azide and 0.5% BSA.

## Données de validation sélectionnées



1X10^6 human PBMCs were surface co-stained with CL750 Anti-Human CD3 and 5 ul PE Anti-Human CD19 (PE-65110, Clone: HIB19) or Mouse IgG1 Isotype Control. Cells were not fixed. Lymphocytes were gated.



1X10^6 human PBMCs were surface stained with 5 ul PE Anti-Human CD19 (PE-65110, Clone: HIB19) (red) or Mouse IgG1 Isotype Control. Cells were not fixed. Lymphocytes were gated.