À des fins de recherche uniquement

Anticorps Monoclonal anti-CD27

Numéro de catalogue:PE-65132



Informations de base

Numéro de catalogue:

Numéro d'acquisition GenBank: BC012160 Méthode de purification:

PE-65132

DC012100

Purification par affinité

Taille:

Identification du gène (NCBI):

CloneNo.: 0323

100tests , 5 µl/test Hôte:

Nom complet:

Excitation/Emission maxima

Mouse

CD27 molecule MW calculé wavelengths:

Isotype: IgG1, kappa

29 kDa

496 nm, 565 nm / 578 nm

Applications

Applications testées:

FC

Spécificité de l'espèce:

Humain

Informations générales

CD27 (also known as TNFRSF7) is a type I glycoprotein expressed on some B cells and the majority of T cells, and is a member of the tumor necrosis factor (TNF) receptor family. CD27 is required for generation and long-term maintenance of T cell immunity (PMID: 11062504). It is a receptor for CD70 (CD27L). Ligation of CD27 by CD70 induces strong ubiquitination of TRAF and the activation of both canonical and non-canonical NF-kappaB pathways, as well as the JNK pathway (PMID: 19426224). CD27 may also play a role in apoptosis through association with SIVA1.

Stockage

Stockage:


Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

Tampon de stockage:


PBS with 0.09% sodium azide and 0.5% BSA.

W: ptglab.com

Données de validation sélectionnées



1X10^6 human PBMCs were surface co-stained with APC Anti-Human CD3 and 5 ul PE Anti-Human CD27 (PE-65132, Clone:O323) or Mouse IgG1 Isotype Control. Cells were not fixed. Lymphocytes were gated.



1X10^6 human PBMCs were surface stained with 5 ul PE Anti-Human CD27 (PE-65132, Clone:0323) or Mouse IgG1 Isotype Control. Cells were not fixed. Lymphocytes were gated.