

À des fins de recherche uniquement

Anticorps Monoclonal anti-CD19

Numéro de catalogue: PE-65145



Informations de base

Numéro de catalogue:

PE-65145

Taille:

100 tests, 5 µl/test

Hôte:

Mouse

Isotype:

IgG1, kappa

Numéro d'acquisition GenBank:

BC006338

Identification du gène (NCBI):

930

Nom complet:

CD19 molecule

MW calculé

556 aa, 61 kDa

Méthode de purification:

Purification par affinité

CloneNo.:

SJ25C1

Excitation/Emission maxima
wavelengths:

496 nm, 565 nm / 578 nm

Applications

Applications testées:

FC

Spécificité de l'espèce:

Humain

Informations générales

CD19 is a 95 kDa type I transmembrane glycoprotein belonging to the immunoglobulin superfamily (PMID: 2472450). It is expressed by B cells and follicular dendritic cells. CD19 is up-regulated at the step of B-lineage commitment during the differentiation of the hematopoietic stem cell, it remains on during subsequent stages of differentiation until finally down-regulated during terminal differentiation into plasma cells (PMID: 8528044). CD19 is involved in B cell development, activation and differentiation. It is the dominant component for the signaling complex on B cells that includes CD21 (CR2), CD81 (TAPA-1) and CD225 and acts as a critical co-receptor for BCR signal transduction (PMID: 23210908).

Stockage

Stockage:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

Tampon de stockage:

Tampon à base de phosphate avec azoture de sodium à 0,09 % et gélatine à 0,1 %, pH 7,2.



For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



1X10⁶ human peripheral blood lymphocytes were surface stained with 0.25 ug PE-anti-human CD19 (PE-65145, clone SJ25C1) (red) or 0.25 ug PE-mouse IgG1 isotype control (blue). Cells were not fixed.

0.1 mL human whole blood cells were surface stained with 0.25 ug APC-anti-human CD3 (APC-65151, clone UCHT1) and 0.25 ug PE-anti-human CD19 (PE-65145, clone SJ25C1). Cells were then treated with red blood cell lysis buffer and were gated for lymphocytes for analysis of CD3 and CD19 staining. Cells were not fixed.