

À des fins de recherche uniquement

Anticorps Monoclonal anti-CD73

Numéro de catalogue: PE-65162



Informations de base

Numéro de catalogue:	BC015940	Méthode de purification:
PE-65162	L'anticorps purifié est conjugué à la R-phycoérythrine (PE) dans des conditions optimales. Le conjugué est purifié par chromatographie d'exclusion de taille.	
Taille:	Identification du gène (NCBI):	CloneNo.:
100 tests, 5 µl/test	4907	AD2
Hôte:	Nom complet:	Excitation/Emission maxima wavelengths:
Mouse	5'-nucleotidase, ecto (CD73)	496 nm, 565 nm / 578 nm
Isotype:	MW calculé	
IgG1, kappa	29 kDa, 63 kDa	

Applications

Applications testées:
FC

Spécificité de l'espèce:
Humain

Informations générales

CD73, also known as ecto-5'-nucleotidase (5'-NT), is a 70-kDa, glycosyl-phosphatidylinositol-linked membrane-bound glycoprotein found in most tissues (PMID: 18404475; 20179192). CD73 is an ectoenzyme that catalyzes the dephosphorylation of AMP and other nucleoside monophosphates (PMID: 9553767). In the human immune system, CD73 is expressed on subsets of T and B cells, on germinal center follicular dendritic cells, and on thymic medullary reticular fibroblasts and epithelial cells (PMID: 2137649; 9553767). CD73 is highly expressed in many human solid tumors and is closely involved in cancer progression (PMID: 20179192).


Stockage

Stockage:
Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.
Tampon de stockage:
PBS with 0.09% sodium azide and 0.5% BSA.


For technical support and original validation data for this product please contact:
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



1×10^6 human PBMCs were surface co-stained with APC Anti-Human CD3 and 5 μ l PE Anti-Human CD73 (PE-65162, Clone:AD2) or Mouse IgG1 Isotype Control. Cells were not fixed. Lymphocytes were gated.



1×10^6 human PBMCs were surface stained with 5 μ l PE Anti-Human CD73 (PE-65162, Clone:AD2) or Mouse IgG1 Isotype Control. Cells were not fixed. Lymphocytes were gated.