

For Research Use Only

PYK2 Polyclonal antibody

Catalog Number: 10044-2-AP **1 Publications**



Basic Information

Catalog Number: 10044-2-AP	GenBank Accession Number: BC036651	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 200 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 2185	Recommended Dilutions: WB 1:500-1:1000 IHC 1:20-1:200
Source: Rabbit	Full Name: PTK2B protein tyrosine kinase 2 beta	
Isotype: IgG	Calculated MW: 120 kDa	
Immunogen Catalog Number: AG0002	Observed MW: 116 kDa	

Applications

Tested Applications: IHC, WB, ELISA	Positive Controls:
Cited Applications: WB	WB : mouse brain tissue, Jurkat cells
Species Specificity: human, mouse	IHC : human medulloblastoma tissue, human kidney tissue
Cited Species: rat	

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

Proline-rich tyrosine kinase 2 (Pyk2; also known as CAK, RAFTK and CADTK) is a cytoplasmic tyrosine kinase implicated to play a role in several intracellular signaling pathways. It is expressed in many cells and tissues and migrated as a 130 kDa band in Western Blotting analysis.

Notable Publications

Author	Pubmed ID	Journal	Application
L Yang	28983639	Herz	WB

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

***** 20ul sizes contain 0.1% BSA**


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



mouse brain tissue were subjected to SDS PAGE followed by western blot with 10044-2-AP (PYK2 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human medulloblastoma using 10044-2-AP (PYK2 antibody) at dilution of 1:50 (under 10x lens).