

Progranulin/PGRN Polyclonal antibody

Catalog Number: 10053-1-AP

2 Publications

Basic Information

Catalog Number:	GenBank Accession Number:	Purification Method:
10053-1-AP	BC010577	Antigen affinity purification
Size:	GeneID (NCBI):	Recommended Dilutions:
150ul, Concentration: 1500 µg/ml by Nanodrop;	2896	WB 1:500-1:1000 IHC 1:50-1:500
Source:	Full Name:	
Rabbit	granulin	
Isotype:	Calculated MW:	
IgG	64 kDa	
Immunogen Catalog Number:	Observed MW:	
AG0010	64 kDa	

Applications

Tested Applications:	Positive Controls:
IHC, WB, ELISA	WB: A431 cells, HEK-293 cells, MCF-7 cells
Cited Applications:	IHC: human colon cancer tissue, human liver cancer tissue
IF, IHC	
Species Specificity:	
human	
Cited Species:	
human	

Note-IHC: suggested antigen retrieval with **TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with **citrate buffer pH 6.0****

Background Information

GRN, also known as PGRN or PCDGF, is a cysteine-rich protein of 68.5 kDa that is typically secreted into a highly glycosylated 88 kDa form. PGRN is a unique growth factor that plays an important role in cutaneous wound healing. It has an anti-inflammatory effect and promotes cell proliferation. When PCDGF is degraded to several 6-25 kDa fragments, called granulins (GRNs) by neutrophil proteases, a pro-inflammatory reaction occurs. PGRN is widely expressed, particularly in epithelial cells, immune cells, neurons, and chondrocytes. High levels of PGRN expression have been reported in human cancers, and its expression is closely correlated with the development and metastasis of several cancers. The recent discovery that mutations in the gene encoding for pro-granulin (GRN) cause frontotemporal lobar degeneration (FTLD), and other neurodegenerative diseases leading to dementia, has brought renewed interest in progranulin and its functions in the central nervous system.

Notable Publications

Author	Pubmed ID	Journal	Application
Guanshen Cui	37084418	Cell Prolif	IF
Chen Xiang-yu XY	18706200	Chin Med J (Engl)	IHC

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
 T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
 E: proteintech@ptglab.com
 W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 10053-1-AP (Granulin antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human colon cancer tissue slide using 10053-1-AP (Granulin antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human colon cancer tissue slide using 10053-1-AP (Granulin antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).