

Nectin 2 Polyclonal antibody

Catalog Number: 10100-2-AP

1 Publications

Basic Information

Catalog Number: 10100-2-AP	GenBank Accession Number: BC003091	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 133 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 5819	Recommended Dilutions: WB 1:200-1:1000
Source: Rabbit	Full Name: poliovirus receptor-related 2 (herpesvirus entry mediator B)	
Isotype: IgG	Calculated MW: 58 kDa	
Immunogen Catalog Number: AG0149		

Applications

Tested Applications: FC, WB, ELISA	Positive Controls:
Cited Applications: IF, WB	WB : mouse kidney tissue, 3T3-L1 cells, mouse heart tissue
Species Specificity: human, mouse	
Cited Species: monkey	

Background Information

Poliovirus receptor-related 2 (herpesvirus entry mediator B) (PVRL2, synonyms: HVEB, PRR2, CD112, PVRR2) is an adhesion molecule widely expressed in cell lines of different lineages, including hematopoietic, neuronal, endothelial and epithelial cells. CD112 belongs to a new family of immunoglobulin-like molecules that includes four members (CD111, CD112, PRR3 and CD155) sharing an ectodomain made of three Ig domains, of V, C, C types. CD112 is expressed in the myelo-monocytic and megakaryocytic hematopoietic lineages and the function in hematopoiesis is currently unknown. CD112 is an intercellular homophilic adhesion. CD112 localizes specifically at adherens junctions via its cytoplasmic interaction with the scaffold F-actin binding protein afadin. This interaction is mediated by a sequence located to the C terminal ends of both isoforms of CD112 (A/ExYV) and the PDZ domain of afadin. This sequence is also found in CD112 and PRR3 and represents a specific consensus in the family, which is also named nectin family. Disruption of the murine CD112 gene leads to infertility of male mice with morphologically aberrant spermatozoa. CD112 mediates entry of some alphaherpesvirus mutants (also named HveB) via its V domain. CD112 is involved in cell to cell spreading of viruses

Notable Publications

Author	Pubmed ID	Journal	Application
Dudak Amanda A	21371776	Eur J Cell Biol	WB, IF

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



mouse kidney tissue were subjected to SDS PAGE followed by western blot with 10100-2-AP (Nectin 2 antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.



1X10⁶ MCF-7 cells were stained with 0.2ug Nectin 2 antibody (10100-2-AP, red) and control antibody (blue). Fixed with 4% PFA blocked with 3% BSA (30 min). Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) with dilution 1:1500.