#### For Research Use Only

# VGLL1 Polyclonal antibody

Catalog Number:10124-2-AP

**Featured Product** 

4 Publications



**Basic Information** 

Catalog Number: GenBank Accession Number:

10124-2-AP BC000045
Size: GeneID (NCBI):
150ul , Concentration: 1000 µg/ml by 51442
Nanodrop: LANDROT LD:

Nanodrop; UNIPROT ID:
Source: Q99990
Rabbit Full Name:

Isotype: vestigial like 1 (Drosophila)

IgG Calculated MW:
Immunogen Catalog Number: 29 kDa
AG0169 Observed MW:

26-29 kDa

Purification Method: Antigen affinity purification Recommended Dilutions: WB 1:1000-1:6000 IHC 1:20-1:200

**Applications** 

Tested Applications: IHC, WB, ELISA Cited Applications: WB,RIP,IHC,IF

Species Specificity: human, mouse, rat Cited Species: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

#### Positive Controls:

WB: Jurkat cells, human brain tissue, rat brain tissue, PC-3 cells, mouse brain tissue, mouse spleen tissue, rat spleen tissue

IHC: human prostate cancer tissue,

### **Background Information**

VGLL1 is a specific coactivator for the mammalian TEFs. It can bind proteins of the TEA domain family of transcription factors. Vgll1-TEFs complex upregulates the expression of IGFBP-5, a proliferation-promoting gene, and facilitates anchorage-independent cell proliferation. The calculated molecular weight of VGLL1 is 29 kDa, and the observed molecular weight is 26-29 kDa(PMID:31816819,33087697).

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
María Ángeles Castilla	24891455	Endocr Relat Cancer	
Mi-Aie Hwang	36613819	Int J Mol Sci	WB
Bo-Kyung Kim	31816819	Cancers (Basel)	WB,IHC

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA

## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 10124-2-AP (VGLL1 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human prostate cancer using 10124-2-AP (VGLL1 antibody) at dilution of 1:50 (under 10x lens).