

For Research Use Only

GNAS Polyclonal antibody

Catalog Number: 10150-2-AP

Featured Product

7 Publications



Basic Information

Catalog Number: 10150-2-AP	GenBank Accession Number: BC002722	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 500 µg/ml by Nanodrop;	GeneID (NCBI): 2778	Recommended Dilutions: WB 1:2000-1:12000 IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate IHC 1:200-1:800 IF 1:50-1:500
Source: Rabbit	Full Name: GNAS complex locus	
Isotype: IgG	Calculated MW: 45 kDa	
Immunogen Catalog Number: AG0197	Observed MW: 46 kDa	

Applications

Tested Applications: IF, IHC, IP, WB, ELISA	Positive Controls:
Cited Applications: IF, IHC, WB	WB : Jurkat cells, mouse brain tissue, HepG2 cells, MCF-7 cells, mouse liver tissue, rat liver tissue, rat brain tissue
Species Specificity: human, mouse, rat	IP : mouse liver tissue,
Cited Species: human, mouse	IHC : mouse brain tissue, human pancreas tissue, human colon tissue
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	IF : HeLa cells,

Background Information

Guanine nucleotide binding protein (G protein), alpha stimulating activity polypeptide 1 (GNAS1) is the ubiquitously expressed heterotrimeric G protein that couples receptors to the effector enzyme adenylyl cyclase and is required for receptor-stimulated intracellular cAMP generation. Mutations of Gs(alpha) residues involved in the GTPase reaction that lead to constitutive activation are present in endocrine tumors, fibrous dysplasia of bone, and McCune-Albright syndrome. The molecular weight of Gs(alpha) protein is about 46 kDa.

Notable Publications

Author	Pubmed ID	Journal	Application
Gemoll Timo T	21290163	Cell Mol Life Sci	WB
Hongda Ding	32123532	Cell Mol Biol Lett	WB,IF
Lele Liu	32036696	Am J Physiol Renal Physiol	WB,IF

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 10150-2-AP (GNAS antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours.




HepG2 cells were subjected to SDS PAGE followed by western blot with 10150-2-AP (GNAS antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



IP Result of anti-GNAS (IP:10150-2-AP, 4ug; Detection:10150-2-AP 1:300) with mouse liver tissue lysate 4000ug.



Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 10150-2-AP (GNAS antibody) at dilution of 1:400 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 10150-2-AP (GNAS antibody) at dilution of 1:400 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using GNAS antibody (10150-2-AP) at dilution of 1:200 and CoraLite@488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), 594-phalloidine (red).