## For Research Use Only

# CDC42 Polyclonal antibody

Catalog Number: 10155-1-AP

Featured Product

**54 Publications** 

BC002711



**Basic Information** 

Catalog Number: GenBank Accession Number:

ze: Genel D (NCBI):

150ul , Concentration: 600 µg/ml by 998

Nanodrop; Full Name:

Source: cell division cycle 42 (GTP binding

Rabbit protein, 25kDa)
Isotype: Calculated MW:
IgG 21 kDa
Immunogen Catalog Number: Observed MW:

AG0207 21 kDa

**Applications** 

Tested Applications:

IF, IHC, WB, ELISA

10155-1-AP

Cited Applications: CoIP, IF, IHC, WB

Species Specificity:

human, mouse, rat

Cited Species:

human, chicken, rat, mouse, hamster, bovine

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Purification Method:

Antigen affinity purification

**Recommended Dilutions:** 

WB 1:500-1:2000 IHC 1:50-1:500 IF 1:20-1:200

Positive Controls:

WB: HT-29 cells, HeLa cells, C6 cells, Jurkat cells,

NIH/3T3 cells, mouse brain tissue

IHC: human pancreas cancer tissue, human pancreas

tissue, human skeletal muscle tissue

IF: C6 cells, HeLa cells

# **Background Information**

Cdc42, the human homologue of the cell division cycle 42 of Saccharomyces cerevisiae, belongs to the Rho/Rac subfamily of Ras-like GTPases. It acts as molecular switches to regulate numerous cellular responses including the organization of the actin cytoskeleton, gene transcription, cell cycle progression, and membrane trafficking. The molecular weight is about 21 kd.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Yu-Yuan Zhang	36259516	J Biol Chem	WB
Xiang Pan	28979695	Am J Transl Res	WB
Teng Zhou	25212524	Biomaterials	WB

## Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


# **Selected Validation Data**




Various lysates were subjected to SDS PAGE followed by western blot with 10155-1-AP (CDC 42 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human pancreas cancer tissue slide using 10155-1-AP (CDC42 Antibody) at dilution of 1:200 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human pancreas cancer tissue slide using 10155-1-AP (CDC42 Antibody) at dilution of 1:200 (under 40x lens).



C6 cells were subjected to SDS PAGE followed by western blot with 10155-1-AP (CDC42 Antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Ethanol) fixed C6 cells using 10155-1-AP (CDC 42 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).