For Research Use Only

PPARD Polyclonal antibody

Catalog Number: 10156-2-AP

23 Publications



Purification Method:

WB 1:500-1:1000

IHC 1:10-1:100

Antigen affinity purification

Recommended Dilutions:

Basic Information

Catalog Number: GenBank Accession Number: 10156-2-AP BC002715

Size: GeneID (NCBI):

150ul, Concentration: 450 µg/ml by Nanodrop and 153 µg/ml by Bradford Full Name:

method using BSA as the standard;

receptor delta Rabbit Calculated MW: 50 kDa Isotype:

54 kDa Immunogen Catalog Number:

AG0208

IgG

Positive Controls:

WB: mouse heart tissue, mouse skeletal muscle tissue,

peroxisome proliferator-activated

Observed MW:

COLO 320 cells

IHC: human ovary tumor tissue,

Applications

Tested Applications: IHC, WB, ELISA **Cited Applications:** IF, IHC, WB

Species Specificity: human, mouse, rat **Cited Species:** human, rat, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

Peroxisome proliferator-activator receptor delta(PPARD), a ligand-activated transcription factor, belongs to the PPARs family, which get its name for their chemicals that induce proliferation of peroxisomes, organelles that contributes to the oxidation of fatty acids. PPARD is a receptor for peroxisome proliferator and once activated by a ligand, it binds to and activates the promoter elements of target genes, such as acylCoA oxidase gene. But it act as a repressor for the NPC1L1 gene.

Notable Publications

Author	Pubmed ID	Journal	Application
Juan Ji	36142731	Int J Mol Sci	WB
Juan Ji	27789312	Neuropharmacology	WB
Yaqiong Zhang	27815817	Neurotox Res	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

*** 20ul sizes contain 0.1% BSA

Aliquoting is unnecessary for -20°C storage


in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com W: ptglab.com

Selected Validation Data



mouse heart tissue were subjected to SDS PAGE followed by western blot with 10156-2-AP (PPARD antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human ovary tumor using 10156-2-AP (PPARD antibody) at dilution of 1:10 (under 10x lens).