For Research Use Only

## ATG5 Polyclonal antibody

Catalog Number:10181-2-AP

Featured Product






Basic Information	Catalog Number: 10181-2-AP	GenBank Accession Number: BC002699	Purification Method: Antigen affinity purification	
	Size: 150ul , Concentration: 950 ug/ml by Nanodrop; Source:	GenelD (NCBI): 9474	Recommended Dilutions: WB 1:1000-1:5000	
		UNIPROT ID: Q9H1Y0	IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate	
	Rabbit Isotype: IgG	Full Name: ATG5 autophagy related 5 homolog (S. cerevisiae)		
	Immunogen Catalog Number: AG0214	Calculated MW: 32 kDa		
		Observed MW: 32 kDa, 40-45 kDa, 50-55 kDa		
Applications	Tested Applications:	Positive Controls:		
	WB, IP, ELISA Cited Applications:		WB : A549 cells, HeLa cells, mouse kidney tissue, HepG2 cells	
	WB, IHC, IF, IP, CoIP Species Specificity: human, mouse, rat	IP : HeLa cells,		
	Cited Species: human, mouse, rat, pig, monkey, chi	ken, bovine,		
	hamster, goat, grouper			
Background Information	ATG5, also named as APG5L and ASP role in the apoptotic process, possibl the apoptotic process, occurring dow autophagosomic-lysosomal degrada ubiquitin-like conjugation system in vesicles. It mediates autophagosom	y within the modified cytoskeletc nstream of caspase activity. Autop tion of bulk cytoplasmic contents which Atg12 is covalently bound e-independent host protection. Th e the ATG5-ATG12 complex (55 k	required for autophagy. It plays an importar on. Its expression is a relatively late event ir phagy is a catabolic process for the . Formation of the autophagosome involves to Atg5 and targeted to autophagosome is antibody is raised against 28-275 amino Da) which can be truncated and generate a 4	
	ATG5, also named as APG5L and ASP role in the apoptotic process, possibl the apoptotic process, occurring dow autophagosomic-lysosomal degrada ubiquitin-like conjugation system in vesicles. It mediates autophagosom acids of human ATG5. It can recogniz 45 kDa band. 10181-2-AP also recogn	y within the modified cytoskeletc nstream of caspase activity. Autop tion of bulk cytoplasmic contents which Atg12 is covalently bound e-independent host protection. Th e the ATG5-ATG12 complex (55 k	m. Its expression is a relatively late event in phagy is a catabolic process for the Formation of the autophagosome involves to Atg5 and targeted to autophagosome is antibody is raised against 28-275 amino	
	ATG5, also named as APG5L and ASP role in the apoptotic process, possibl the apoptotic process, occurring dow autophagosomic-lysosomal degrada ubiquitin-like conjugation system in vesicles. It mediates autophagosom acids of human ATG5. It can recogniz 45 kDa band. 10181-2-AP also recogn	y within the modified cytoskeleto hstream of caspase activity. Autop tion of bulk cytoplasmic contents which Atg12 is covalently bound a-independent host protection. Th e the ATG5-ATG12 complex (55 k hizes the free ATG5 (32 kDa).	m. Its expression is a relatively late event in phagy is a catabolic process for the Formation of the autophagosome involves to Atg5 and targeted to autophagosome is antibody is raised against 28-275 amino Da) which can be truncated and generate a 4	
	ATG5, also named as APG5L and ASP role in the apoptotic process, possibl the apoptotic process, occurring dow autophagosomic-lysosomal degrada ubiquitin-like conjugation system in vesicles. It mediates autophagosom acids of human ATG5. It can recogniz 45 kDa band. 10181-2-AP also recogn Author Pul Zeen Zhu 362	y within the modified cytoskeleto hstream of caspase activity. Autop tion of bulk cytoplasmic contents which Atg12 is covalently bound e-independent host protection. Th e the ATG5-ATG12 complex (55 k hizes the free ATG5 (32 kDa).	n. Its expression is a relatively late event in phagy is a catabolic process for the . Formation of the autophagosome involves to Atg5 and targeted to autophagosome is antibody is raised against 28-275 amino Da) which can be truncated and generate a 2 Application	
Background Information	ATG5, also named as APG5L and ASP role in the apoptotic process, possibil the apoptotic process, occurring dow autophagosomic-lysosomal degrada ubiquitin-like conjugation system in vesicles. It mediates autophagosom acids of human ATG5. It can recogniz 45 kDa band. 10181-2-AP also recogn Author Pul Zeen Zhu 362 Samana Batool 302	y within the modified cytoskeleto hstream of caspase activity. Autop tion of bulk cytoplasmic contents which Atg12 is covalently bound e-independent host protection. Th e the ATG5-ATG12 complex (55 k nizes the free ATG5 (32 kDa). med ID Journal 248959 Front Oncol	n. Its expression is a relatively late event in phagy is a catabolic process for the Formation of the autophagosome involves to Atg5 and targeted to autophagosome is antibody is raised against 28-275 amino Da) which can be truncated and generate a Application WB	


For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free<br/>in USA), or 1(312) 455-8498 (outside USA)E: proteintech@ptglab.comW: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data




HeLa cells were subjected to SDS PAGE followed by western blot with 10181-2-AP (ATG5 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



WB result of ATG5 antibody (10181-2-AP; 1:2000; room temperature for 1.5 hours) with wild-type and ATG5 knockout HeLa cells.



Various lysates were subjected to SDS PAGE followed by western blot with 10181-2-AP (ATG5 antibody) at dilution of 1:2500 incubated at room temperature for 1.5 hours.



IP result of anti-ATG5 (IP:10181-2-AP, 4ug; Detection:10181-2-AP 1:3000) with HeLa cells lysate 1760 ug.