For Research Use Only

Caspase 6/P18/P11 Polyclonal antibody

Catalog Number:10198-1-AP

Featured Product

13 Publications




Basic Information	Catalog Number: 10198-1-AP	GenBank Accession Nun BC000305		Method: inity purification			
	Size:	GeneID (NCBI):	C	Recommended Dilutions: WB 1:500-1:1000 IHC 1:50-1:500			
	150ul , Concentration: 450 ug/ml by	839					
	Nanodrop and 293 ug/ml by Bradford	UNIPROT ID:	IHC 1:50-1:				
	method using BSA as the standard;	P55212					
	Source: Rabbit	Full Name: caspase 6, apoptosis-related cysteine peptidase Calculated MW: 33 kDa, 22 kDa					
	Isotype: IgG Immunogen Catalog Number: AG0257						
					Observed MW: 33-35 kDa		
					Applications	11 A A A A A A A A A A A A A A A A A A	
		WB, IHC, ELISA		WB: Staurosporine treated Jurkat cells, Jurkat cells,			
Cited Applications: WB, IF		nouse brain tissue					
Species Specificity: human, mouse, rat		IHC : human lung cancer tissue, human prostate canc tissue, human skin tissue					
Cited Species: human, mouse, rat							
Note-IHC: suggested antigen r TE buffer pH 9.0; (*) Alternativ retrieval may be performed w buffer pH 6.0	vely, antigen						
Background Information	Caspase-6 belongs to caspase family of cysteinyl-aspartate specific proteases.Precursor of CASP6 produces two subunits, large (18kDa) and small (16kDa) that dimerize. It cleaves poly(ADP-ribose) polymerase, as well as lamins and is involved in the activation cascade of caspases responsible for apoptosis execution. Researches showed that CASP6 could be an early instigator of neuronal dysfunction and regulates B cell activation and differentiation into plasma cells by modifying cell cycle entry. IRAK3 is an important target for CASP6. It can reveal five bands of 28, 32 36, 49, and 64 kDa in human neurons and fetal brain in western blot, the 32 and 28 kDa bands represent procaspase-6 is more abundant than pro-arm caspase-6 is not abundant than pro-arm caspase-6 is more abundant than pro-arm caspase						
Notable Publications	Author Pub	med ID Journal		Application			
			iol Interact	WB			
	Weixin Hou 345	26765 Drug De	s Devel Ther	WB			
	Danli Lu 361	81629 Transl S	troke Res	WB			
Storage	Storage: Store at -20°C. Stable for one year after Storage Buffer:						
	PBS with 0.02% sodium azide and 50° Aliquoting is unnecessary for -20°C st	•••					

in USA), or 1(312) 455-8498 (outside USA)

ב. proterntecn@ptg W: ptglab.com

ıy other manufacturer.


Selected Validation Data




Staurosporine treat Jurkat cells were subjected to SDS PAGE followed by western blot with 10198-1-AP (Caspase 6/p18/p11 antibody at dilution of 1:600 incubated at room temperature for 1.5 hours.




Jurkat cells were subjected to SDS PAGE followed by western blot with 10198-1-AP (CASP6 antibody) at dilution of 1:800 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human prostate cancer using 10198-1-AP (CASP6 antibody) at dilution of 1:50 (under 10x lens).



WB result of Caspase 6/p18/p11 antibody (10198-1-AP; 1:600; incubated at room temperature for 1.5 hours) with sh-Control and sh-Caspase 6/p18/p11 transfected Jurkat cells.



Immunohistochemical analysis of paraffinembedded human lung cancer using 10198-1-AP (CASP6 antibody) at dilution of 1:200 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human lung cancer using 10198-1-AP (CASP6 antibody) at dilution of 1:200 (under 40x lens).