

NCK2 Polyclonal antibody

Catalog Number: 10206-1-AP

1 Publications

Basic Information

Catalog Number: 10206-1-AP	GenBank Accession Number: BC000103	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 600 µg/ml by Nanodrop and 300 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 8440	Recommended Dilutions: WB 1:500-1:1000 IHC 1:50-1:500
Source: Rabbit	Full Name: NCK adaptor protein 2	
Isotype: IgG	Calculated MW: 43 kDa	
Immunogen Catalog Number: AG0262	Observed MW: 43 kDa	

Applications

Tested Applications: IHC, WB, ELISA	Positive Controls:
Cited Applications: WB	WB : U-937 cells,
Species Specificity: human, mouse, rat	IHC : human placenta tissue,
Cited Species: mouse	

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

Lots of the protein-protein interactions that are essential for eukaryotic intracellular signal transduction are mediated by protein binding modules including SH2, SH3, and LIM domains. NCK is a SH3- and SH2-containing adaptor protein implicated in coordinating various signaling pathways, including those of growth factor receptors and cell adhesion receptors. NCK-2 is a ubiquitously expressed and NCK-related adaptor protein comprising primarily three N-terminal SH3 domains and one C-terminal SH2 domain. NCK-2 interacts with PINCH, a LIM-only protein implicated in integrin-linked kinase signaling, and recognizes several key components of growth factor kinase-signaling pathways including EGF receptors, PDGF receptor, and IRS-1. NCK-2 may thus function as an adaptor protein connecting the growth factor receptor-signaling pathways with the integrin-signaling pathways. NCK-2 also interacts with focal adhesion kinase (FAK), a cytoplasmic protein tyrosine kinase critically involved in the cellular control of motility.

Notable Publications

Author	Pubmed ID	Journal	Application
Mi Huang	32508946	Evid Based Complement Alternat Med	WB

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



U-937 cells were subjected to SDS PAGE followed by western blot with 10206-1-AP (NCK2 Antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human placenta tissue slide using 10206-1-AP (NCK2 Antibody) at dilution of 1:200 (under 10x lens).



Immunohistochemical analysis of paraffin-embedded human placenta tissue slide using 10206-1-AP (NCK2 Antibody) at dilution of 1:200 (under 40x lens).