

For Research Use Only

NCK2 Polyclonal antibody

Catalog Number: 10206-1-AP

Featured Product

2 Publications



Basic Information

Catalog Number:

10206-1-AP

Size:

150ul, Concentration: 600 ug/ml by Nanodrop and 300 ug/ml by Bradford method using BSA as the standard;

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG0262

GenBank Accession Number:

BC000103

GeneID (NCBI):

8440

UNIPROT ID:

O43639

Full Name:

NCK adaptor protein 2

Calculated MW:

43 kDa

Observed MW:

43 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:1000

IHC 1:50-1:500

Applications

Tested Applications:

WB, IHC, ELISA

Cited Applications:

WB

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse

Positive Controls:

WB : U-937 cells,

IHC : human placenta tissue,

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

Lots of the protein-protein interactions that are essential for eukaryotic intracellular signal transduction are mediated by protein binding modules including SH2, SH3, and LIM domains. NCK is a SH3- and SH2-containing adaptor protein implicated in coordinating various signaling pathways, including those of growth factor receptors and cell adhesion receptors. NCK-2 is a ubiquitously expressed and NCK-related adaptor protein comprising primarily three N-terminal SH3 domains and one C-terminal SH2 domain. NCK-2 interacts with PINCH, a LIM-only protein implicated in integrin-linked kinase signaling, and recognizes several key components of growth factor kinase-signaling pathways including EGF receptors, PDGF receptor, and IRS-1. NCK-2 may thus function as an adaptor protein connecting the growth factor receptor-signaling pathways with the integrin-signaling pathways. NCK-2 also interacts with focal adhesion kinase (FAK), a cytoplasmic protein tyrosine kinase critically involved in the cellular control of motility.

Notable Publications

Author	Pubmed ID	Journal	Application
Mi Huang	32508946	Evid Based Complement Alternat Med	WB
Xiaodong Li	38583810	Biochem Pharmacol	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



U-937 cells were subjected to SDS PAGE followed by western blot with 10206-1-AP (NCK2 Antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human placenta tissue slide using 10206-1-AP (NCK2 Antibody) at dilution of 1:200 (under 10x lens).



Immunohistochemical analysis of paraffin-embedded human placenta tissue slide using 10206-1-AP (NCK2 Antibody) at dilution of 1:200 (under 40x lens).