For Research Use Only

Chemerin/RARRES2 Polyclonal antibody

Catalog Number: 10216-1-AP

Featured Product

7 Publications

GenBank Accession Number:



Basic Information

Catalog Number:

10216-1-AP BC000069
Size: GeneID (NCBI):

150ul , Concentration: 650 ug/ml by 5919 Nanodrop and 333 ug/ml by Bradford UNIPROT ID:

Nanodrop and 333 ug/ml by Bradfor method using BSA as the standard;

Source: Full Name:

Rabbit retinoic acid receptor responder

Isotype: (tazarotene induced) 2

Q99969

sotype: (tazarotene induced) 2 gG Calculated MW:

Immunogen Catalog Number: 163 aa, 19 kDa
AG0261 Observed MW:
14-19 kDa

Applications

Tested Applications:

WB, IHC, ELISA

Cited Applications:

WB, IHC, IF

Species Specificity:

human, mouse

Cited Species:

human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:200-1:1000 IHC 1:50-1:500

Positive Controls:

WB: human plasma tissue,

IHC: human placenta tissue, mouse liver tissue

Background Information

Chemerin, also named as TIG2 and RARRES2, is adipocyte-secreted protein (adipokine) that regulates adipogenesis, metabolism and inflammation through activation of the chemokine-like receptor 1 (CMKLR1). Its other ligands include G protein-coupled receptor 1 (GPR1) and chemokine receptor-like 2 (CCRL2). It positively regulates adipocyte differentiation, modulates the expression of adipocyte genes involved in lipid and glucose metabolism and might play a role in angiogenesis, a process essential for the expansion of white adipose tissue. RARRES2 have both pro- and anti-inflammatory properties depending on the modality of enzymatic cleavage by different classes of proteases. The MW of Chemerin is 14-19 kDa.

Notable Publications

Author	Pubmed ID	Journal	Application
Ming Yu	36523492	Front Pharmacol	WB,IHC
Xiaoyuan Hu	35155249	Front Oncol	IHC
Xuan Zhou	33843164	J Zhejiang Univ Sci B	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffe

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


*** 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:


T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com W: ptglab.com This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Immunohistochemical analysis of paraffinembedded human placenta tissue slide using 10216-1-AP (Chemerin antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human placenta tissue slide using 10216-1-AP (Chemerin antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



human plasma were subjected to SDS PAGE followed by western blot with 10216-1-AP (Chemerin antibody at dilution of 1:300 incubated at room temperature for 1.5 hours.