For Research Use Only

PDLIM7 Polyclonal antibody

Catalog Number:10221-1-AP

Featured Product 14 Publications




Basic Information	Catalog Number: 10221-1-AP	GenBank Accession Num BC001093	ber: Purification Method: Antigen affinity purification
	Size: GeneID (NCBI):		Recommended Dilutions:
	150ul , Concentration: 800 ug/ml by		
	Nanodrop and 400 ug/ml by Bradford	UNIPROT ID:	IP 0.5-4.0 ug for 1.0-3.0 mg of total
	method using BSA as the standard;	Q9NR12	proteinlysate
	Source:	Full Name:	IHC 1:200-1:800 IF/ICC 1:50-1:500
	Rabbit	Rabbit PDZ and LIM domain 7 (enigma)	
	lsotype: Calculated MW: IgG 55/60 kDa		
	Immunogen Catalog Number: AG0264	Observed MW: 55 kDa	
Applications	Tested Applications:	Positive Controls:	
			VB : SKOV-3 cells, HeLa cells, MCF-7 cells
	Cited Applications: WB, IHC, IF, IP, CoIP	II	P : MCF-7 cells,
		П	HC : human colon tissue, human ovary tumor tissue
	Species Specificity: human	П	F/ICC : SKOV-3 cells,
	Cited Species:		
	human, mouse		
	TE buffer pH 9.0; (*) Alternativ retrieval may be performed w buffer pH 6.0		
	bujjer pri 0.0		
Background Information	PDLIM7 (PDZ and LIM domain protein of PDZ-LIM proteins which function as C-terminal LIM domain. Its LIM doma	scaffold to interact with ins bind to proteins involv ell invasion and metastas	multiple proteins through N-terminal PDZ domain a /ed in mitogenic signaling, and it plays a role in sis. In addition, PDLIM7 is an essential positive
	PDLIM7 (PDZ and LIM domain protein of PDZ-LIM proteins which function as C-terminal LIM domain. Its LIM doma tumorigenesis by promoting cancer of regulator of osteoblast differentiation	scaffold to interact with ins bind to proteins involv ell invasion and metastas	multiple proteins through N-terminal PDZ domain a /ed in mitogenic signaling, and it plays a role in sis. In addition, PDLIM7 is an essential positive
	PDLIM7 (PDZ and LIM domain protein of PDZ-LIM proteins which function as C-terminal LIM domain. Its LIM domai tumorigenesis by promoting cancer or regulator of osteoblast differentiation Author Pub	scaffold to interact with r ins bind to proteins involv ell invasion and metastas n and maturation and bon	multiple proteins through N-terminal PDZ domain a ved in mitogenic signaling, and it plays a role in sis. In addition, PDLIM7 is an essential positive e formation. Application
Background Information	PDLIM7 (PDZ and LIM domain protein of PDZ-LIM proteins which function as C-terminal LIM domain. Its LIM domai tumorigenesis by promoting cancer of regulator of osteoblast differentiation Author Pub Michael D Healy 363	scaffold to interact with in ins bind to proteins involved in and maturation and bone med ID Journal	sis. In addition, PDLIM7 is an essential positive e formation. Application WB
	PDLIM7 (PDZ and LIM domain protein of PDZ-LIM proteins which function as C-terminal LIM domain. Its LIM domain tumorigenesis by promoting cancer of regulator of osteoblast differentiation Author Pub Michael D Healy 363 Hui Liu 253	scaffold to interact with in ins bind to proteins involved invasion and metastas n and maturation and bon med ID Journal 02387 Structure	multiple proteins through N-terminal PDZ domain a ved in mitogenic signaling, and it plays a role in isis. In addition, PDLIM7 is an essential positive e formation. Application WB Res WB
	PDLIM7 (PDZ and LIM domain protein of PDZ-LIM proteins which function as C-terminal LIM domain. Its LIM domain tumorigenesis by promoting cancer of regulator of osteoblast differentiation Author Pub Michael D Healy 363 Hui Liu 253	scaffold to interact with n ins bind to proteins involve ell invasion and metastas n and maturation and bon med ID Journal 02387 Structure 36289 J Orthop 42427 Cell Rep er shipment.	multiple proteins through N-terminal PDZ domain a ved in mitogenic signaling, and it plays a role in isis. In addition, PDLIM7 is an essential positive e formation. Application WB Res WB

For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




WB result of PDLIM7 antibody (10221-1-AP; 1:100000; incubated at room temperature for 1.5 hours) with sh-Control and sh-PDLIM7 transfected HeLa cells.



SKOV-3 cells were subjected to SDS PAGE followed by western blot with 10221-1-AP (PDLIM7 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Ethanol) fixed SKOV-3 cells using PDLIM7 antibody (10221-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



IP result of anti-PDLIM7 (IP:10221-1-AP, 4ug: Detection:10221-1-AP 1:20000) with MCF-7 cells lysate 1000 ug.



Immunohistochemical analysis of paraffinembedded human colon tissue slide using 10221-1-AP (PDLIM7 antibody) at dilution of 1:400 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).