

For Research Use Only

CRABP2 Polyclonal antibody

Catalog Number: 10225-1-AP

Featured Product

37 Publications



Basic Information

Catalog Number: 10225-1-AP	GenBank Accession Number: BC001109	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 350 µg/ml by Nanodrop;	GeneID (NCBI): 1382	Recommended Dilutions: WB 1:1000-1:4000 IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate
Source: Rabbit	Full Name: cellular retinoic acid binding protein 2	IHC 1:50-1:500 IF 1:50-1:500
Isotype: IgG	Calculated MW: 16 kDa	
Immunogen Catalog Number: AG0309	Observed MW: 16 kDa	

Applications

Tested Applications: FC (Intra), IF, IHC, IP, WB, ELISA	Positive Controls:
Cited Applications: CoIP, IF, IHC, IP, WB	WB : HEK-293 cells, HeLa cells, HT-29 cells, MCF-7 cells, mouse embryo tissue
Species Specificity: human, mouse, rat	IP : HeLa cells,
Cited Species: human, mouse	IHC : human skin tissue,
	IF : MCF-7 cells, human skin cancer tissue
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	

Background Information

Cellular retinoic acid binding protein 2 (CRABP2, synonyms: RBP6, CRABP-II). A number of specific carrier proteins for members of the vitamin A family have been discovered. Cellular retinoic acid binding proteins (CRABP) are low molecular weight proteins whose precise function remains unknown. CRABP2 is important in retinoic acid-mediated regulation of human skin growth and differentiation. It has been postulated that the CRABP2 gene is transcriptionally regulated by a newly synthesized regulatory protein.

Notable Publications

Author	Pubmed ID	Journal	Application
Masanori Goto	26348989	Brain Res	IF
Yasuhiro Adachi	36089341	J UOEH	WB
Xin Liu	31736873	Front Endocrinol (Lausanne)	WB,IHC

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Immunohistochemical analysis of paraffin-embedded human skin using 10225-1-AP (CRABP2 antibody) at dilution of 1:50 (under 10x lens).




Immunofluorescent analysis of (4% PFA) fixed MCF-7 cells using CRABP2 antibody (10225-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-phalloidin (red).




WB result of CRABP2 antibody (10225-1-AP; 1:300; incubated at room temperature for 1.5 hours) with sh-Control and sh-CRABP2 transfected HeLa cells.



IP Result of anti-CRABP2 (IP:10225-1-AP, 3ug; Detection:10225-1-AP 1:300) with HeLa cells lysate 4650ug.



Various lysates were subjected to SDS PAGE followed by western blot with 10225-1-AP (CRABP2 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



1X10⁶ MCF-7 cells were intracellularly stained with 0.2 ug Anti-Human CRABP2 (10225-1-AP) and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.2 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).