

For Research Use Only

# S100A11 Polyclonal antibody

Catalog Number: 10237-1-AP

Featured Product

51 Publications



## Basic Information

### Catalog Number:

10237-1-AP

### Size:

150ul, Concentration: 550 ug/ml by Nanodrop and 460 ug/ml by Bradford method using BSA as the standard;

### Source:

Rabbit

### Isotype:

IgG

### Immunogen Catalog Number:

AG0343

### GenBank Accession Number:

BC001410

### GeneID (NCBI):

6282

### UNIPROT ID:

P31949

### Full Name:

S100 calcium binding protein A11

### Calculated MW:

105 aa, 12 kDa

### Observed MW:

12 kDa

### Purification Method:

Antigen affinity purification

### Recommended Dilutions:

WB 1:2000-1:12000

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

IHC 1:50-1:500

IF/ICC 1:200-1:800

## Applications

### Tested Applications:

WB, IHC, IF/ICC, FC (Intra), IP, ELISA

### Cited Applications:

WB, IHC, IF, IP, CoIP, Neutralization

### Species Specificity:

human, rat

### Cited Species:

human, mouse, rat

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

### Positive Controls:

**WB:** DU 145 cells, HCT 116 cells, BxPC-3 cells, human lung tissue, HEK-293 cells, human heart tissue, COLO 320 cells, HaCaT cells, U2OS cells, SKOV-3 cells

**IP:** DU 145 cells, PC-3 cells

**IHC:** human breast cancer tissue, human cervical cancer tissue, human colon tissue, human colon cancer tissue, human liver cancer tissue, human lung cancer tissue, human ovary tumor tissue, human thyroid cancer tissue, mouse kidney tissue, rat kidney tissue

**IF/ICC:** PC-3 cells, BxPC-3 cells

## Background Information

S100A11 (also known as S100C or calziggarin), is a member of the S100 family of proteins containing 2 EF-hand calcium-binding motifs (PMID: 18694925). S100 proteins are localized in the cytoplasm and/or nucleus of a wide range of cells, and involved in the regulation of a number of cellular processes such as cell cycle progression and differentiation. First discovered in 1989, S100A11 has since been proposed to have direct biological functions in an assortment of physiological processes such as endo- and exocytosis, regulation of enzyme activity, cell growth and regulation, apoptosis and inflammation (PMID: 15241500). Chromosomal rearrangements and altered expression of S100A11 have been implicated in tumor metastasis. This S100A11 antibody (10237-1-AP) has also been instrumental in investigations into S100A11's role in the development of brain tumors in tuberous sclerosis complex (TSC) patients (PMID: 20133820).

## Notable Publications

Author	Pubmed ID	Journal	Application
Yu-Pu Liu	33132818	Front Neurosci	IF
Arseniy E Yuzhalin	31545917	Am J Physiol Gastrointest Liver Physiol	IF
Arsila P. K. Ashraf	36200035	Front Cell Dev Biol	IF

## Storage

### Storage:

Store at -20°C. Stable for one year after shipment.

### Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 10237-1-AP (S100A11 antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours.




Various lysates were subjected to SDS PAGE followed by western blot with 10237-1-AP (S100A11 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




WB result of S100A11 antibody (10237-1-AP, 1:4000) with si-control and si-S100A11 transfected PC-3 cells.




IP result of anti-S100A11 (IP:10237-1-AP, 3ug; Detection:10237-1-AP 1:1000) with PC-3 cells lysate 1000ug.




IP result of anti-S100A11 (IP:10237-1-AP, 4ug; Detection:10237-1-AP 1:5000) with DU 145 cells lysate 1280 ug.




Immunohistochemical analysis of paraffin-embedded human liver cancer tissue slide using 10237-1-AP (S100A11 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).






Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 10237-1-AP (S100A11 Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded human cervical cancer tissue slide using 10237-1-AP (S100A11 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded human lung cancer tissue slide using 10237-1-AP (S100A11 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human thyroid cancer tissue slide using 10237-1-AP (S100A11 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 10237-1-AP (S100A11 Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




1x10<sup>6</sup> PC-3 cells were intracellularly stained with 0.8 ug S100A11 Polyclonal antibody (10237-1-AP) and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2)(red), or 0.8 ug Rabbit IgG control Rabbit PolyAb (30000-O-AP) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).

Immunohistochemical analysis of paraffin-embedded mouse kidney tissue slide using 10237-1-AP (S100A11 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunofluorescent analysis of (-20°C Ethanol) fixed PC-3 cells using S100A11 antibody (10237-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).

Immunohistochemical analysis of paraffin-embedded rat kidney tissue slide using 10237-1-AP (S100A11 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed BxPC-3 cells using S100A11 antibody (10237-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red).



Immunohistochemical analysis of paraffin-embedded human colon tissue slide using 10237-1-AP (S100A11 antibody) at dilution of 1:1200 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).