

For Research Use Only

S100A11 Polyclonal antibody, PBS Only

Catalog Number: 10237-1-PBS

Featured Product



Basic Information

Catalog Number: 10237-1-PBS	GenBank Accession Number: BC001410	Purification Method: Antigen affinity purification
Size: 100ug, Concentration: 1 mg/ml by Nanodrop;	GeneID (NCBI): 6282	
Source: Rabbit	UNIPROT ID: P31949	
Isotype: IgG	Full Name: S100 calcium binding protein A11	
Immunogen Catalog Number: AG0343	Calculated MW: 105 aa, 12 kDa	
	Observed MW: 12 kDa	

Applications

Tested Applications:
WB, IHC, IF/ICC, FC (Intra), IP, ELISA

Species Specificity:
human, mouse, rat

Background Information

S100A11 (also known as S100C or calziggarin), is a member of the S100 family of proteins containing 2 EF-hand calcium-binding motifs (PMID: 18694925). S100 proteins are localized in the cytoplasm and/or nucleus of a wide range of cells, and involved in the regulation of a number of cellular processes such as cell cycle progression and differentiation. First discovered in 1989, S100A11 has since been proposed to have direct biological functions in an assortment of physiological processes such as endo- and exocytosis, regulation of enzyme activity, cell growth and regulation, apoptosis and inflammation (PMID: 15241500). Chromosomal rearrangements and altered expression of S100A11 have been implicated in tumor metastasis. This S100A11 antibody (10237-1-AP) has also been instrumental in investigations into S100A11's role in the development of brain tumors in tuberous sclerosis complex (TSC) patients (PMID: 20133820).

Storage

Storage:
Store at -80°C.

Storage Buffer:
PBS only, pH7.3


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 10237-1-AP (S100A11 antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 10237-1-PBS in a different storage buffer formulation.




WB result of S100A11 antibody (10237-1-AP, 1:4000) with si-control and si-S100A11 transfected PC-3 cells. This data was developed using the same antibody clone with 10237-1-PBS in a different storage buffer formulation.




Various lysates were subjected to SDS PAGE followed by western blot with 10237-1-AP (S100A11 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 10237-1-PBS in a different storage buffer formulation.




IP result of anti-S100A11 (IP:10237-1-AP, 3ug; Detection:10237-1-AP 1:1000) with PC-3 cells lysate 1000ug. This data was developed using the same antibody clone with 10237-1-PBS in a different storage buffer formulation.




IP result of anti-S100A11 (IP:10237-1-AP, 4ug; Detection:10237-1-AP 1:5000) with DU 145 cells lysate 1280 ug. This data was developed using the same antibody clone with 10237-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffin-embedded human liver cancer tissue slide using 10237-1-AP (S100A11 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 10237-1-PBS in a different storage buffer formulation.






Immunohistochemical analysis of paraffin-embedded human cervical cancer tissue slide using 10237-1-AP (S100A11 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 10237-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffin-embedded human lung cancer tissue slide using 10237-1-AP (S100A11 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 10237-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffin-embedded human thyroid cancer tissue slide using 10237-1-AP (S100A11 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 10237-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffin-embedded mouse kidney tissue slide using 10237-1-AP (S100A11 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 10237-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffin-embedded rat kidney tissue slide using 10237-1-AP (S100A11 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 10237-1-PBS in a different storage buffer formulation.





Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 10237-1-AP (S100A11 Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 10237-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 10237-1-AP (S100A11 Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 10237-1-PBS in a different storage buffer formulation.

Immunofluorescent analysis of (-20°C Ethanol) fixed PC-3 cells using S100A11 antibody (10237-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L). This data was developed using the same antibody clone with 10237-1-PBS in a different storage buffer formulation.

Immunofluorescent analysis of (-20°C Ethanol) fixed BxPC-3 cells using S100A11 antibody (10237-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red). This data was developed using the same antibody clone with 10237-1-PBS in a different storage buffer formulation.



1x10⁶ PC-3 cells were intracellularly stained with 0.8 ug S100A11 Polyclonal antibody (10237-1-AP) and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2)(red), or 0.8 ug rabbit IgG isotype control (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer. This data was developed using the same antibody clone with 10237-1-PBS in a different storage buffer formulation.

Immunohistochemical analysis of paraffin-embedded human colon tissue slide using 10237-1-AP (S100A11 antibody) at dilution of 1:1200 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 10237-1-PBS in a different storage buffer formulation.