

For Research Use Only

HDAC3 Polyclonal antibody

Catalog Number: 10255-1-AP

Featured Product

62 Publications



Basic Information

Catalog Number:

10255-1-AP

Size:

150ul, Concentration: 500 ug/ml by Nanodrop;

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG0396

GenBank Accession Number:

BC000614

GeneID (NCBI):

8841

UNIPROT ID:

O15379

Full Name:

histone deacetylase 3

Calculated MW:

49 kDa

Observed MW:

49 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:3000

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

IHC 1:20-1:200

IF/ICC 1:50-1:500

Applications

Tested Applications:

WB, IHC, IF/ICC, IP, ELISA

Cited Applications:

WB, IHC, IF, IP, CoIP, ChIP

Species Specificity:

human

Cited Species:

human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : A431 cells, HL-60 cells, HeLa cells, U-251 cells, HEK-293 cells, HepG2 cells, Jurkat cells, K-562 cells

IP : A431 cells,

IHC : human ovary tumor tissue,

IF/ICC : K-562 cells,

Background Information

Histone acetylation/deacetylation alters chromosome structure and affects transcription factor access to DNA. Histone deacetylase (HDAC) and histone acetyltransferase (HAT) are enzymes that regulate transcription by selectively deacetylating or acetylating the (-amino groups of lysines located near the amino termini of core histone proteins. At least 4 classes of HDAC were identified. HDAC3 is a class I HDAC. HDAC3 has histone deacetylase activity and may participate in the regulation of transcription through its binding with the zinc-finger transcription factor YY1. HDAC3 can also down-regulate p53 function and thus modulate cell growth and apoptosis. The gene encoding HDAC3 is regarded as a potential tumor suppressor gene. This antibody is a rabbit polyclonal antibody raised against an internal region of human HDAC3.

Notable Publications

| Author | Pubmed ID | Journal | Application |
|--------------|-----------|--------------------|-------------|
| Rui Han | 36175803 | Med Oncol | WB |
| Zhengrong Gu | 36106389 | J Clin Lab Anal | IHC |
| Hairui Yuan | 25213552 | Acta Physiol (Oxf) | WB |

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 10255-1-AP (HDAC3 antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.




WB result of HDAC3 antibody (10255-1-AP; 1:3000; incubated at room temperature for 1.5 hours) with sh-Control and sh-HDAC3 transfected U-251 cells.




Immunohistochemical analysis of paraffin-embedded human ovary tumor using 10255-1-AP (HDAC3 antibody) at dilution of 1:50 (under 10x lens).



Immunohistochemical analysis of paraffin-embedded human ovary tumor using 10255-1-AP (HDAC3 antibody) at dilution of 1:50 (under 40x lens).



IP result of anti-HDAC3 (IP:10255-1-AP, 3ug; Detection:10255-1-AP 1:300) with A431 cells lysate 1200ug.



Immunofluorescent analysis of (4% PFA) fixed K-562 cells using HDAC3 antibody (10255-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).