

For Research Use Only

IFNGR2 Polyclonal antibody

Catalog Number: 10266-1-AP **7 Publications**



Basic Information

Catalog Number: 10266-1-AP	GenBank Accession Number: BC003624	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 260 µg/ml by Nanodrop and 150 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 3460	Recommended Dilutions: WB 1:500-1:1000 IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate IHC 1:50-1:500
Source: Rabbit	UNIPROT ID: P38484	
Isotype: IgG	Full Name: interferon gamma receptor 2 (interferon gamma transducer 1)	
Immunogen Catalog Number: AG0317	Calculated MW: 45 kDa	
	Observed MW: 50-55 kDa	

Applications

Tested Applications: WB, IP, FC, IHC, ELISA	Positive Controls: WB : A549 cells, HeLa cells, HepG2 cells IP : A549 cells, IHC : human breast cancer tissue,
Cited Applications: WB, IF	
Species Specificity: human	
Cited Species: human, mouse	

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

Interferon-gamma (IFN- γ) is a cytokine critical for innate and adaptive immunity against viral and intracellular bacterial infections and for tumor control. Cellular responses to IFN- γ are activated through its interaction with a heterodimeric receptor consisting of two subunits, IFNGR1 and IFNGR2. IFNGR1 is the ligand-binding subunit which binds IFN- γ with high affinity, whereas IFNGR2 serves as the accessory subunit which attaches to IFN- γ with a significantly lower affinity. Two subunits bind one IFN- γ dimer. Defects in IFNGR2 are a cause of mendelian susceptibility to mycobacterial disease (MSMD), also known as familial disseminated atypical mycobacterial infection. (PMID: 17981204; 946248; 10888113)

Notable Publications

Author	Pubmed ID	Journal	Application
Kefang Tan	28562647	PLoS One	WB
Timothy J Break	33446526	Science	WB
Qingnan Wu	34347390	Cancer Commun (Lond)	WB

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




A549 cells were subjected to SDS PAGE followed by western blot with 10266-1-AP (IFNGR2 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 10266-1-AP (IFNGR2 antibody) at dilution of 1:200 (under 10x Lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 10266-1-AP (IFNGR2 antibody) at dilution of 1:200 (under 40x Lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-IFNGR2 (IP:10266-1-AP, 4ug; Detection:10266-1-AP 1:500) with A549 cells lysate 1800ug.



1X10⁶ HeLa cells were stained with 0.2ug IFNGR2 antibody (10266-1-AP, red) and control antibody (blue). Fixed with 90% MeOH blocked with 3% BSA (30 min). Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) with dilution 1:1500.