### For Research Use Only

# PAX8 Polyclonal antibody

Catalog Number: 10336-1-AP

Featured Product

527 Publications



#### **Basic Information**

Catalog Number:

10336-1-AP

Size:

150ul , Concentration: 500 ug/ml by

Nanodrop;

Source: Rabbit

Isotype:

GenBank Accession Number:

BC001060

GeneID (NCBI):

UNIPROT ID:

Q06710

Full Name: paired box 8

Calculated MW:

48 kDa
Observed MW:

48 kDa, 58 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:2000-1:10000 IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:800-1:3200 IF-P 1:200-1:800 IF/ICC 1:200-1:800

# **Applications**

**Tested Applications:** 

WB, IHC, IF/ICC, IF-P, IP, ELISA

Cited Applications:

WB, IHC, IF, CoIP, ELISA

Species Specificity:

human, mouse

Cited Species:

human, mouse, rat, pig, rabbit, zebrafish, bovine

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

#### **Positive Controls:**

WB: SKOV-3 cells, HEK-293 cells, Raji cells, NIH/3T3

cells

IP: SKOV-3 cells, HeLa cells

IHC: human ovary cancer tissue, human renal cell carcinoma tissue, human thyroid cancer tissue, mouse

kidney tissue IF-P: mouse kidney tissue,

IF/ICC : SKOV-3 cells.

# **Background Information**

PAX8 is a member of the paired box (PAX) family of transcription factors, typically containing a paired box domain and a paired-type homeodomain. It is expressed during organogenesis of the thyroid gland, kidney and Mullerian system. It is thought to regulate the expression of Wilms tumor suppressor (WT1) gene and mutations in PAX8 have been associated with Wilms tumor cells, thyroid and ovarian carcinomas. PAX8 is a useful marker in distinguishing ovarian carcinomas from mammary carcinomas (PMID: 18724243). PAX8 is expressed in a high percentage of ovarian serous, endometrioid, and clear cell carcinomas, but only rarely in primary ovarian mucinous adenocarcinomas.

This antibody can detect all 5 isoforms of PAX8 (31 kDa, 34 kDa, 41 kDa, 43 kDa and 48 kDa). A 58-60 kDa form has also been reported (PMID:15650356, 1049952). Proteintech's 10336-1-AP antibody has cross-reactivities with the other PAX family members.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Yong Kyun Kim	28905451	Stem Cells	IF
Olga Kondrashova	30266954	Nat Commun	IHC
Bilhan	27765277	Curr Res Transl Med	IHC

# Storage

Storage

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


\*\*\* 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:


T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com W: ptglab.com This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


# **Selected Validation Data**




SKOV-3 cells were subjected to SDS PAGE followed by western blot with 10336-1-AP (PAX8 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.




Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded mouse kidney tissue using PAX8 antibody (10336-1-AP) at dilution of 1:400 and Multi-rAb CoraLite ® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0)



Immunofluorescent analysis of (4% PFA) fixed SKOV-3 cells using PAX8 antibody (10336-1-AP) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-phalloidin (red).



IP result of anti-PAX8 (IP:10336-1-AP, 4ug; Detection:10336-1-AP 1:15000) with SKOV-3 cells lysate 3280 ug.



Immunohistochemical analysis of paraffinembedded human ovarian cancer slide using 10336-1-AP (PAX8 antibody) at dilution of 1:1600 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).