For Research Use Only

RNH1 Polyclonal antibody

Catalog Number: 10345-1-AP

Featured Product

11 Publications



Basic Information

Catalog Number: GenBank Accession Number:

10345-1-AP BC000677 GeneID (NCBI):

150ul, Concentration: 700 µg/ml by 6050

Nanodrop; Full Name:

Source: ribonuclease/angiogenin inhibitor 1

Rabbit Calculated MW: Isotype: 50 kDa

IgG Observed MW: Immunogen Catalog Number: 45-50 kDa

AG0350

Applications

Tested Applications:

FC, IF, IHC, IP, WB, ELISA Cited Applications:

IF, IHC, WB Species Specificity:

human, mouse, rat **Cited Species:**

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

human, mouse

buffer pH 6.0

Purification Method:

Antigen affinity purification

Recommended Dilutions: WB 1:2000-1:10000

IP 0.5-4.0 ug for IP and 1:500-1:1000

for WB IHC 1:50-1:500 IF 1:200-1:800

Positive Controls:

WB: HEK-293T cells, HeLa cells, human brain tissue, human heart tissue, human liver tissue, LO2 cells, HepG2 cells, Jurkat cells, human placenta tissue IP: mouse liver tissue, HEK-293T cells, rat liver tissue

IHC: mouse liver tissue, IF: MCF-7 cells,

Background Information

Ribonuclease/ angiogenin inhibitor (RNH, synonyms: RAI, MGC 4569, MGC 18200, MGC 54054) is a protein that binds tightly to ribonucleases in cells and may be essential in the control of mRNA degradation and gene expression. The human RNH gene has been regionally localized to chromosome band 11p15 by in situ hybridization.

Notable Publications

Author	Pubmed ID	Journal	Application
Yasutoshi Akiyama	36361884	Int J Mol Sci	WB
Yasutoshi Akiyama	35300117	Front Mol Biosci	WB
Shozo Honda	26124144	Proc Natl Acad Sci U S A	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 10345-1-AP (RNH1 antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours.




WB result of RNH1 antibody (10345-1-AP, 1:2500) with si-control and si-RNH1 transfected HeLa cells.




Immunohistochemical analysis of paraffinembedded mouse liver tissue slide using 10345-1-AP (RNH1 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded mouse liver tissue slide using 10345-1-AP (RNH1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunofluorescent analysis of (-20°C Methanol) fixed MCF-7 cells using RNH1 antibody (10345-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).




IP Result of anti-RNH1 (IP:10345-1-AP, 4ug; Detection:10345-1-AP 1:500) with mouse liver tissue lysate 6400ug.




IP result of anti-RNH1(IP:10345-1-AP, 4ug; Detection:10345-1-AP 1:8000) with HEK-293T cells lysate 2040 ug.



IP result of anti-RNH1(IP:10345-1-AP, 4ug; Detection:10345-1-AP 1:8000) with mouse liver tissue lysate 1840 ug.



IP result of anti-RNH1(IP:10345-1-AP, 4ug; Detection:10345-1-AP 1:8000) with rat liver tissue lysate 5520 ug.



1X10^6 MCF-7 cells were intracellularly stained with 0.4 ug Anti-Human RNH1 (10345-1-AP) and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).