For Research Use Only

BCAS3 Polyclonal antibody

Catalog Number: 10402-1-AP

Featured Product



Basic Information

Catalog Number: GenBank Accession Number:

10402-1-AP Antigen affinity purification BC001250 GeneID (NCBI): Size: Recommended Dilutions: WB 1:500-1:2000

150ul, Concentration: 170 µg/ml by 54828 Nanodrop and 133 µg/ml by Bradford Full Name:

method using BSA as the standard;

breast carcinoma amplified sequence

Rabbit Calculated MW: 99 kDa Isotype: IgG Observed MW: Immunogen Catalog Number: 74 kDa

AG0646

Applications Tested Applications:

WB, ELISA

Species Specificity: human, mouse, rat

Positive Controls:

WB: A549 cells, A431 cells, HeLa cells, Jurkat cells

Purification Method:

Background Information

Rudhira/BCAS3 (Breast Cancer Amplified Sequence 3) is a conserved protein expressed in the embryonic vasculature and malignant tumors. Rudhira is a predicted WD40 domain protein. During mouse development Rudhira is expressed in erythropoietic and angiogenic precursors but not in their differentiated progeny. This transient expression of Rudhira is also seen in normal adult angiogenesis. BCAS3, the human ortholog of Rudhira is 98% identical and the gene maps to a breakpoint of hematological neoplasms on chromosome 17q23.1. BCAS3 is also expressed in vascular development in embryoid bodies in vitro and was originally identified as a gene overexpressed in breast cancers. It is reported to be a target of metastasis associated protein-1 (MTA-1). Further BCAS3 expression is also upregulated in grade III and IV glioblastomas where it is misexpressed in the tumor cells.

Storage


Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data





Various lysates were subjected to SDS PAGE followed by western blot with 10402-1-AP (BCAS3 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.

WB result of BCAS3 antibody (10402-1-AP; 1:600; incubated at room temperature for 1.5 hours) with sh-Control and sh-BCAS3 transfected A431 cells.