For Research Use Only

DMAP1 Polyclonal antibody

Catalog Number: 10411-1-AP

Featured Product

6 Publications



Basic Information

Catalog Number:

GenBank Accession Number:

Antigen affinity purification

10411-1-AP Size:

GeneID (NCBI):

Recommended Dilutions:

150ul, Concentration: 400 µg/ml by

55929

BC002855

WB 1:1000-1:4000

for WB

Purification Method:

Nanodrop and 293 µg/ml by Bradford Full Name:

DNA methyltransferase 1 associated

IP 0.5-4.0 ug for IP and 1:200-1:1000

method using BSA as the standard;

Isotype:

Calculated MW: 53 kDa

IgG Immunogen Catalog Number:

Observed MW:

AG0665

Rabbit

53-60 kDa

Applications

Tested Applications:

IP, WB, ELISA

Cited Applications:

ChIP, RIP, WB

Species Specificity:

human, mouse, rat

Cited Species: human, mouse **Positive Controls:**

WB: COLO 320 cells, HeLa cells, NIH/3T3 cells

IP: COLO 320 cells,

Background Information

DMAP1 (DNA methyltransferase 1-associated protein 1) a component of the NuA4 histone acetyltransferase complex, was originally identified as an interacting molecule with DNA methyltransferase 1 (DNMT1) which colocalizes with PCNA at DNA replication foci. DMAP1 is required for DNA repair and acts against genomic instability. DMAP1 is widely implicated in transcription repression and activation via its interaction with multiple targets including histone deacetylase2 (HDAC2). The antibody reacts specifically with DMAP1 protein.

Notable Publications

Author	Pubmed ID	Journal	Application
Yue Ming	36327957	Psychiatry Investig	ChIP,RIP
Jinbeom Heo	31709755	EMBO Mol Med	WB
Poshen B Chen	26551076	Nat Struct Mol Biol	

Storage

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 10411-1-AP (DMAP1 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



NIH/3T3 cells were subjected to SDS PAGE followed by western blot with 10411-1-AP (DMAP1 Antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



IP Result of anti-DMAP1 (IP:10411-1-AP, 3ug; Detection:10411-1-AP 1:300) with COLO 320 cells lysate 2500ug.