

BAD Polyclonal antibody

Catalog Number: 10435-1-AP

93 Publications

Basic Information

Catalog Number:

10435-1-AP

Size:

150ul, Concentration: 387 µg/ml by Nanodrop and 387 µg/ml by Bradford method using BSA as the standard;

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG0688

GenBank Accession Number:

BC001901

GeneID (NCBI):

572

UNIPROT ID:

Q92934

Full Name:

BCL2-associated agonist of cell death

Calculated MW:

18 kDa

Observed MW:

18-25 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:3000

IHC 1:100-1:500

Applications

Tested Applications:

WB, IHC, FC (Intra), ELISA

Cited Applications:

WB, IHC

Species Specificity:

human, mouse

Cited Species:

human, mouse, rat, pig, sheep

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : HeLa cells, Jurkat cells, HepG2 cells, Neuro-2a cells

IHC : human prostate hyperplasia tissue, human prostate cancer tissue, mouse liver tissue, human lymphoma tissue

Background Information

BAD, also named as BBC6 and BCL2L8, promotes cell death. It successfully competes for the binding to Bcl-X(L), Bcl-2 and Bcl-W, thereby affecting the level of heterodimerization of these proteins with BAX. BAD can reverse the death repressor activity of Bcl-X(L), but not that of Bcl-2. It appears to act as a link between growth factor receptor signaling and the apoptotic pathways.

Notable Publications

Author	Pubmed ID	Journal	Application
Yanan Li	27663303	BMC Microbiol	WB
Xuan Zhou	36150222	Chemosphere	WB
Minjing Li	31546831	Biomolecules	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 10435-1-AP (BAD antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.




Various lysates were subjected to SDS PAGE followed by western blot with 10435-1-AP (BAD antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



1X10⁶ Jurkat cells were intracellularly stained with 0.4 ug Anti-Human BAD (10435-1-AP) and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Isotype Control. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Immunohistochemical analysis of paraffin-embedded human prostate hyperplasia tissue slide using 10435-1-AP (BAD Antibody) at dilution of 1:200 (under 40x lens).



Immunohistochemical analysis of paraffin-embedded human prostate hyperplasia tissue slide using 10435-1-AP (BAD Antibody) at dilution of 1:200 (under 10x lens).