For Research Use Only

## HTR3A Polyclonal antibody

Catalog Number:10443-1-AP

Featured Product

6 Publications




Basic Information	Catalog Number: 10443-1-AP	GenBank Accession Number: BC002354	Purification Method: Antigen affinity purification	
	Size: 150ul , Concentration: 260 µg/ml by Nanodrop and 133 µg/ml by Bradford method using BSA as the standard;	GenelD (NCBI):	Recommended Dilutions: WB 1:500-1:1000	
		3359		
		UNIPROT ID: P46098	IHC 1:20-1:200	
	Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG0289	Full Name:		
		5-hydroxytryptamine (serotonin) receptor 3A		
		Calculated MW:		
		55 kDa		
		Observed MW: 55 kDa		
Applications	Tested Applications:	Positive Controls:		
			ouse lung tissue, human brain tissue	
	Cited Applications: WB, IF, IHC	IHC : human small intestine tissue, human prostate cancer tissue		
	Species Specificity: human, mouse Cited Species: human, rat, mouse			
				Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0
			ith citrate	
Background Information	buffer pH 6.0 HTR3A (5-hydroxytryptamine receptor is the subunit A of the type 3 receptor as a neurotransmitter, a hormone, and identified. HTR3A can form functiona	or 3A, 5-HT3A) belongs to the liganc for 5-hydroxytryptamine (5-HT, se d a mitogen. So far, five distinct 5-H l homomers or heteromers with HTI		
	buffer pH 6.0 HTR3A (5-hydroxytryptamine receptor is the subunit A of the type 3 receptor as a neurotransmitter, a hormone, and identified. HTR3A can form functiona receptors are capable of mediating fa (PNS). (PMID: 23038271; 17073663)	or 3A, 5-HT3A) belongs to the liganc for 5-hydroxytryptamine (5-HT, se d a mitogen. So far, five distinct 5-F l homomers or heteromers with HTI ist excitatory neurotransmission in	rotonin), a biogenic hormone that function IT3 receptor subunits (A-E) have been R3B or HTR3C or HTR3D or HTR3E. 5-HT3 the CNS and peripheral nervous system	
	buffer pH 6.0 HTR3A (5-hydroxytryptamine receptor is the subunit A of the type 3 receptor as a neurotransmitter, a hormone, and identified. HTR3A can form functiona receptors are capable of mediating fa (PNS). (PMID: 23038271; 17073663) Author Pub	or 3A, 5-HT3A) belongs to the ligand for 5-hydroxytryptamine (5-HT, see d a mitogen. So far, five distinct 5-H l homomers or heteromers with HTI ist excitatory neurotransmission in med ID Journal	rotonin), a biogenic hormone that functio IT3 receptor subunits (A-E) have been R3B or HTR3C or HTR3D or HTR3E. 5-HT3 the CNS and peripheral nervous system Application	
	buffer pH 6.0HTR3A (5-hydroxytryptamine receptor is the subunit A of the type 3 receptor as a neurotransmitter, a hormone, and identified. HTR3A can form functiona receptors are capable of mediating fa (PNS). (PMID: 23038271; 17073663)AuthorPub Wen-Kai BiWen-Kai Bi321	or 3A, 5-HT3A) belongs to the ligand for 5-hydroxytryptamine (5-HT, see d a mitogen. So far, five distinct 5-H l homomers or heteromers with HT ist excitatory neurotransmission in med ID Journal 71529 Biochem Biophys I	rotonin), a biogenic hormone that functio IT3 receptor subunits (A-E) have been R3B or HTR3C or HTR3D or HTR3E. 5-HT3 the CNS and peripheral nervous system Application Res Commun WB	
Background Information	buffer pH 6.0HTR3A (5-hydroxytryptamine receptor is the subunit A of the type 3 receptor as a neurotransmitter, a hormone, and identified. HTR3A can form functiona receptors are capable of mediating fa (PNS). (PMID: 23038271; 17073663)AuthorPub Wen-Kai BiJing Guo358	or 3A, 5-HT3A) belongs to the ligand for 5-hydroxytryptamine (5-HT, see d a mitogen. So far, five distinct 5-H l homomers or heteromers with HTI ist excitatory neurotransmission in med ID Journal	R3B or HTR3C or HTR3D or HTR3E. 5-HT3 the CNS and peripheral nervous system Application Res Commun WB	
	buffer pH 6.0HTR3A (5-hydroxytryptamine receptor is the subunit A of the type 3 receptor as a neurotransmitter, a hormone, and identified. HTR3A can form functiona receptors are capable of mediating fa (PNS). (PMID: 23038271; 17073663)AuthorPub Wen-Kai BiJing Guo358	or 3A, 5-HT3A) belongs to the ligand for 5-hydroxytryptamine (5-HT, sei d a mitogen. So far, five distinct 5-H l homomers or heteromers with HT ist excitatory neurotransmission in med ID Journal 71529 Biochem Biophys I 65309 Front Endocrinol (I	Application Res Commun WB Lausanne) WB	
	buffer pH 6.0HTR3A (5-hydroxytryptamine receptor is the subunit A of the type 3 receptor as a neurotransmitter, a hormone, and identified. HTR3A can form functiona receptors are capable of mediating fa (PNS). (PMID: 23038271; 17073663)AuthorPub Wen-Kai BiJing Guo358	er shipment.	Application Res Commun WB ausanne) WB	
Notable Publications	buffer pH 6.0HTR3A (5-hydroxytryptamine receptor is the subunit A of the type 3 receptor as a neurotransmitter, a hormone, and identified. HTR3A can form functiona receptors are capable of mediating fa (PNS). (PMID: 23038271; 17073663)AuthorPub Wen-Kai BiWen-Kai Bi321Jing Guo358Jian Tang304Storage: Storage Buffer:	er 3A, 5-HT3A) belongs to the ligance for 5-hydroxytryptamine (5-HT, sei d a mitogen. So far, five distinct 5-H l homomers or heteromers with HT isst excitatory neurotransmission in med ID Journal 71529 Biochem Biophys I 65309 Front Endocrinol (I 05756 Oncol Lett er shipment.	Application Res Commun WB Lausanne) WB	
Notable Publications	buffer pH 6.0HTR3A (5-hydroxytryptamine receptor is the subunit A of the type 3 receptor as a neurotransmitter, a hormone, and identified. HTR3A can form functiona receptors are capable of mediating fat (PNS). (PMID: 23038271; 17073663)AuthorPub Wen-Kai BiWen-Kai Bi321Jing Guo358Jian Tang304Storage: Storage Buffer: PBS with 0.02% sodium azide and 50	er 3A, 5-HT3A) belongs to the ligance for 5-hydroxytryptamine (5-HT, sei d a mitogen. So far, five distinct 5-H l homomers or heteromers with HT isst excitatory neurotransmission in med ID Journal 71529 Biochem Biophys I 65309 Front Endocrinol (I 05756 Oncol Lett er shipment.	Application Res Commun WB ausanne) WB	

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



mouse lung tissue were subjected to SDS PAGE followed by western blot with 10443-1-AP (HTR3A antibody) at dilution of 1:800 incubated at room temperature for 1.5 hours. Immunohistochemical analysis of paraffinembedded human small intestine tissue slide using 10443-1-AP (HTR3A Antibody) at dilution of 1:50 (under 40x lens).



Immunohistochemical analysis of paraffinembedded human small intestine tissue slide using 10443-1-AP (HTR3A Antibody) at dilution of 1:50 (under 10x lens).