

For Research Use Only

BLK Polyclonal antibody

Catalog Number: 10510-1-AP **1 Publications**



Basic Information

Catalog Number: 10510-1-AP	GenBank Accession Number: BC007371	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 600 µg/ml by Nanodrop and 333 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 640	Recommended Dilutions: WB 1:500-1:1000 IP 0.5-4.0 ug for IP and 1:500-1:1000 for WB
Source: Rabbit	Full Name: B Lymphoid tyrosine kinase	Calculated MW: 58 kDa
Isotype: IgG	Observed MW: 58 kDa	IHC 1:50-1:500
Immunogen Catalog Number: AG0795		

Applications

Tested Applications: IHC, IP, WB, ELISA	Positive Controls: WB : Jurkat cells, Raji cells, SH-SY5Y cells
Cited Applications: IP, WB	IP : SH-SY5Y cells,
Species Specificity: human, mouse, rat	IHC : human pancreas cancer tissue,
Cited Species: human	
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	

Background Information

Tyrosine-protein kinase Blk (BLK) is typically involved in cell proliferation and differentiation. BLK is a 58 kDa protein with important role in B-cell receptor signaling and B-cell development. BLK also stimulates INS synthesis and secretion in response to glucose and enhances the expression of several pancreatic beta-cell transcription factors.

Notable Publications

Author	Pubmed ID	Journal	Application
Cefan Zhou	36797256	Signal Transduct Target Ther	WB, IP

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

***** 20ul sizes contain 0.1% BSA**


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Jurkat cells were subjected to SDS PAGE followed by western blot with 10510-1-AP (BLK antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human pancreas cancer tissue slide using 10510-1-AP (BLK Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human pancreas cancer tissue slide using 10510-1-AP (BLK Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP Result of anti-BLK (IP:10510-1-AP, 4ug; Detection:10510-1-AP 1:500) with SH-SY5Y cells lysate 1600ug.