

For Research Use Only

# DAD1 Polyclonal antibody

Catalog Number: 10531-1-AP



## Basic Information

<b>Catalog Number:</b> 10531-1-AP	<b>GenBank Accession Number:</b> BC007403	<b>Purification Method:</b> Antigen affinity purification
<b>Size:</b> 150ul , Concentration: 2000 ug/ml by Nanodrop;	<b>GeneID (NCBI):</b> 1603	<b>Recommended Dilutions:</b> WB 1:1000-1:4000
<b>Source:</b> Rabbit	<b>UNIPROT ID:</b> P61803	IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate
<b>Isotype:</b> IgG	<b>Full Name:</b> defender against cell death 1	IHC 1:50-1:500
<b>Immunogen Catalog Number:</b> AG0814	<b>Calculated MW:</b> 12 kDa	
	<b>Observed MW:</b> 16-20 kDa	

## Applications

<b>Tested Applications:</b> WB, IHC, IP, ELISA	<b>Positive Controls:</b>
<b>Species Specificity:</b> human	WB : HEK-293 cells, HepG2 cells
<b>Note-IHC:</b> suggested antigen retrieval with <b>TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0</b>	IP : HepG2 cells,
	IHC : human liver cancer tissue,

## Background Information

DAD1 (defender against cell death 1) is a downstream target of the NFkB survival pathway and exhibits an antiapoptotic function. It is also named as OST2 (Oligosaccharyl transferase2) and catalyzes the transfer of a high mannose oligosaccharide from a lipid-linked oligosaccharide donor to an asparagine residue within an Asn-X-Ser/Thr consensus motif in nascent polypeptide chains. Increased expression of DAD1 has been observed in various tumors.

## Storage


**Storage:**  
Store at -20°C. Stable for one year after shipment.  
**Storage Buffer:**  
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.  
**Aliquoting is unnecessary for -20°C storage**

\*\*\* 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:  
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)  
E: proteintech@ptglab.com  
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 10531-1-AP (DAD1 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



IP result of anti-DAD1 (IP:10531-1-AP, 4ug; Detection:10531-1-AP 1:700) with HepG2 cells lysate 2600 ug.



Immunohistochemical analysis of paraffin-embedded human liver cancer tissue slide using 10531-1-AP (DAD1 Antibody) at dilution of 1:200 (under 10x lens).



Immunohistochemical analysis of paraffin-embedded human liver cancer tissue slide using 10531-1-AP (DAD1 Antibody) at dilution of 1:200 (under 40x lens).