For Research Use Only

RBM15 Polyclonal antibody

Catalog Number: 10587-1-AP

Featured Product

47 Publications



Basic Information

Catalog Number: GenBank Accession Number:

10587-1-AP BC006397

Size: GeneID (NCBI):
150ul , Concentration: 1000 ug/ml by 64783

Nanodrop; UNIPROT ID:
Source: Q96T37
Rabbit Full Name:

Isotype: RNA binding motif protein 15

IgG Calculated MW:
Immunogen Catalog Number: 105 kDa
AG0938 Observed MW:
100-107 kDa

Purification Method: Antigen affinity purification Recommended Dilutions: WB 1:2000-1:16000

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

protein lysate IHC 1:50-1:500 IF/ICC 1:1000-1:4000

Applications

Tested Applications:

WB, IHC, IF/ICC, IP, ELISA

Cited Applications: WB, IHC, IF, IP, CoIP, RIP Species Specificity:

human, mouse, rat Cited Species: human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: SW480 cells, HEK293 cells, HEK-293 cells

IP: HEK-293 cells,

IHC: human breast cancer tissue, IF/ICC: A431 cells, HepG2 cells

Background Information

RNA-binding motif protein 15 (RBM15) belongs to the SPEN protein family, which have repressor function in several signaling pathway, and may bind to RNA through interaction with spliceosome components. It was first described as a 5' translocation partner of the MAL gene in t(1,22)(p13;q13) infant acute megakaryocytic leukemia, and it functions in murine hematopoiesis through modulating Notch-induced transcriptional activation, which play crucial roles in leukemogenesis. Decreasing RBM15 levels with RNA interference could inhibit the growth and proliferation, block the cell cycle, and induce apoptosis in CML cells.

Notable Publications

Author	Pubmed ID	Journal	Application
Vladimir Majerciak	25234929	RNA	WB
Zhenwei Li	34825733	J Clin Lab Anal	WB
Fanhua Kong	36434140	Discov Oncol	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


*** 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:


T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com W: ptglab.com This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



IP result of anti-RBM15 (IP:10587-1-AP, 4ug; Detection:10587-1-AP 1:500) with HEK-293 cells lysate 2600 ug.




Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 10587-1-AP (RBM15 antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




WB result of RBM15 antibody (10587-1-AP, 1:3000) with si-control and si-RBM15 transfected HEK293



HEK-293 cells were subjected to SDS PAGE followed by western blot with 10587-1-AP (RBM15 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.





SW480 cells were subjected to SDS PAGE followed by western blot with 10587-1-AP (RBM15 antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.

Immunofluorescent analysis of (4% PFA) fixed A431 cells using RBM15 antibody (10587-1-AP) at dilution of 1:2000 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red).



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using RBM15 antibody (10587-1-AP) at dilution of 1:1000 and Multi-rAb Coralite ® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002), CL594-phalloidin (red).