

For Research Use Only

# Hsc70 Polyclonal antibody

Catalog Number: 10654-1-AP

Featured Product

76 Publications



## Basic Information

### Catalog Number:

10654-1-AP

### Size:

150ul, Concentration: 400 ug/ml by Nanodrop;

### Source:

Rabbit

### Isotype:

IgG

### Immunogen Catalog Number:

AG1018

### GenBank Accession Number:

BC007276

### GeneID (NCBI):

3312

### UNIPROT ID:

P11142

### Full Name:

heat shock 70kDa protein 8

### Calculated MW:

70 kDa

### Observed MW:

70 kDa

### Purification Method:

Antigen affinity purification

### Recommended Dilutions:

WB 1:2000-1:16000

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

IHC 1:50-1:500

IF/ICC 1:200-1:800

## Applications

### Tested Applications:

WB, IHC, IF/ICC, IP, ELISA

### Cited Applications:

WB, IHC, IF, IP, CoIP

### Species Specificity:

human, mouse, rat

### Cited Species:

human, mouse, rat, pig

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

### Positive Controls:

**WB** : HEK-293 cells, MCF-7 cells, PC-12 cells, HeLa cells, NIH/3T3 cells, U-87 MG cells, mouse brain tissue, rat brain tissue

**IP** : HEK-293 cells,

**IHC** : human ovary tumor tissue, human renal cell carcinoma tissue

**IF/ICC** : HEK-293 cells,

## Background Information

HSPA8 (also known as HSC70) is a member of the HSPA (HSP70) family of heat-shock proteins which are highly conserved chaperons implicated in protein folding, protein refolding, protein transport, and protein targeting. HSPA8 is a constitutively expressed cytosol/nuclear protein able to translocate between cytoplasm and nucleus. Recently it has been reported that HSPA8 can interact with  $\alpha$ -synuclein, the critical pathological protein of Parkinson's disease, indicating its implication in neurodegenerative disease (21832061). In addition, this antibody is most likely able to recognize Hsp70.

## Notable Publications

Author	Pubmed ID	Journal	Application
Zundong Liu	34556812	Oncogene	WB
Yosuke Okamoto	31506297	J Biol Chem	WB, CoIP
Liu Chih-I CI	22885288	J Proteomics	WB

## Storage

### Storage:

Store at -20°C. Stable for one year after shipment.

### Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 10654-1-AP (Hsc70 antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.




IP result of anti-Hsc70 (IP:10654-1-AP, 4ug; Detection:10654-1-AP 1:2000) with HEK-293 cells lysate 1640 ug.




Immunohistochemical analysis of paraffin-embedded human renal cell carcinoma tissue slide using 10654-1-AP (Hsc70 antibody) at dilution of 1:800 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded human renal cell carcinoma tissue slide using 10654-1-AP (Hsc70 antibody) at dilution of 1:800 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Methanol) fixed HEK-293 cells using Hsc70 antibody (10654-1-AP) at dilution of 1:400 and Coralite®488-Conjugated Goat Anti-Rabbit IgG(H+L).



Immunohistochemical analysis of paraffin-embedded human ovary tumor tissue slide using 10654-1-AP (Hsc70 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human ovary tumor tissue slide using 10654-1-AP (Hsc70 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).