For Research Use Only

FBXW4 Polyclonal antibody

Catalog Number: 10657-1-AP



Purification Method:

WB 1:200-1:1000

IHC 1:50-1:500

Antigen affinity purification

Recommended Dilutions:

Basic Information

Catalog Number: GenBank Accession Number: 10657-1-AP BC007380

GeneID (NCBI): Size:

150ul , Concentration: 260 $\mu g/ml$ by Nanodrop and 173 µg/ml by Bradford Full Name:

method using BSA as the standard; F-box and WD repeat domain

containing 4 Rabbit Calculated MW: 46 kDa Isotype: IgG Observed MW: 50 kDa Immunogen Catalog Number:

AG1047

Applications Tested Applications: Positive Controls:

IHC, WB, ELISA WB: mouse lung tissue, mouse liver tissue

Species Specificity: IHC: human pancreas tissue, human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

F-box proteins have been shown to be critical for the ubiquitin-mediated degradation of cellular regulatory proteins, and they are a family of eukaryotic proteins characterized by an approximately 40 amino acid motif. SCF complex, a class of ubiquitin ligases, consists of invariable components, Skp1 and Cullin, and variable components of F-box proteins, which have a primary role in determining substrate specificity. FBXW4, also known as SHFM3, encodes Fbox and WD-40 domain-containing protein 4. Defects in SHFM3 are a cause of split-hand/foot malformation type 3 (SHFM3), an autosomal dominant disorder characterized by hypoplasia/aplasia of the central digits with fusion of the remaining digits.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


in USA), or 1(312) 455-8498 (outside USA)

W: ptglab.com

Selected Validation Data



mouse lung tissue were subjected to SDS PAGE followed by western blot with 10657-1-AP (FBXW4 Antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human pancreas tissue slide using 10657-1-AP (FBXW4 Antibody) at dilution of 1:200 (under 40x lens).