For Research Use Only

MGP Polyclonal antibody Catalog Number: 10734-1-AP Featured Product

Featured Product






Basic Information	Catalog Number: 10734-1-AP	GenBank Accession Number: BC 005272		Purification Method: Antigen affinity purification			
	Size:	GeneID (NCBI): 4256 UNIPROT ID: P08493 Full Name: matrix Gla protein		Recommended Dilutions: IHC 1:50-1:500 IF-P 1:50-1:500			
	150ul , Concentration: 600 ug/ml by Nanodrop; Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG1091						
					Calculated MW:		
					103 aa, 13 kDa		
					Applications	Tested Applications: IHC, IF-P, ELISA	Positive Controls:
		Cited Applications: WB, IHC, IF		intestine ti		n colon cancer tissue, human small ssue, human breast cancer tissue, human tissue, human breast hyperplasia tissue	
		Species Specificity: human, mouse	IF-P : mouse kidney tissue,				
Cited Species: human, mouse, rat Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0							
						TE buffer pH 9.0; (*) Alternati retrieval may be performed w	vely, antigen
Background Information	TE buffer pH 9.0; (*) Alternati retrieval may be performed w buffer pH 6.0 Matrix Gla protein (MGP), a vitamin I Defects in MGP are the cause of Keuto	vely, antigen vith citrate K-dependent prote el syndrome (KS),	which is an autoso				
	TE buffer pH 9.0; (*) Alternati retrieval may be performed w buffer pH 6.0 Matrix Gla protein (MGP), a vitamin I Defects in MGP are the cause of Keute abnormal cartilage calcification, per	vely, antigen vith citrate (-dependent prote el syndrome (KS), ipheral pulmonan	which is an autoso	mal recessive disorder characterized by			
	TE buffer pH 9.0; (*) Alternati retrieval may be performed w buffer pH 6.0Matrix Gla protein (MGP), a vitamin I Defects in MGP are the cause of Keut abnormal cartilage calcification, perAuthorPu	vely, antigen vith citrate K-dependent prote el syndrome (KS), ipheral pulmonar	which is an autoso y stenosis neural h	mal recessive disorder characterized by earing loss and midfacial hypoplasia.			
_	TE buffer pH 9.0; (*) Alternati retrieval may be performed w buffer pH 6.0Matrix Gla protein (MGP), a vitamin I Defects in MGP are the cause of Keuta abnormal cartilage calcification, perAuthorPut Karolina Sterzyńska30	vely, antigen vith citrate K-dependent prote el syndrome (KS), ipheral pulmonan bmed ID	which is an autoso y stenosis neural h Journal	mal recessive disorder characterized by earing loss and midfacial hypoplasia. Application			
_	TE buffer pH 9.0; (*) Alternati retrieval may be performed w buffer pH 6.0Matrix Gla protein (MGP), a vitamin I Defects in MGP are the cause of Keuta abnormal cartilage calcification, perAuthorPut Karolina SterzyńskaXin He32	vely, antigen vith citrate «-dependent prote el syndrome (KS), ipheral pulmonan bmed ID	which is an autoso y stenosis neural h Journal Int J Mol Sci	Application WB,IHC,IF			
Background Information Notable Publications	TE buffer pH 9.0; (*) Alternati retrieval may be performed w buffer pH 6.0Matrix Gla protein (MGP), a vitamin I Defects in MGP are the cause of Keuta abnormal cartilage calcification, perAuthorPut Karolina SterzyńskaXin He32	vely, antigen vith citrate (-dependent prote el syndrome (KS), ipheral pulmonan (bmed ID) 257426 979364 185284 ter shipment.	which is an autoso y stenosis neural h Journal Int J Mol Sci Exp Cell Res Front Oncol	mal recessive disorder characterized by earing loss and midfacial hypoplasia. Application WB,IHC,IF IHC,WB			


For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 10734-1-AP (MGP antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 10734-1-AP (MGP antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded human breast hyperplasia tissue slide using 10734-1-AP (MGP antibody) at dilution of 1:400 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded human breast hyperplasia tissue slide using 10734-1-AP (MGP antibody) at dilution of 1:400 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded mouse liver tissue using MGP antibody (10734-1-AP) at dilution of 1:400 and CoraLite@488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed mouse kidney tissue using MGP antibody (10734-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed mouse kidney tissue using MGP antibody (10734-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).