

For Research Use Only

NF-κB p65 Polyclonal antibody

Catalog Number: 10745-1-AP

Featured Product

1552 Publications



Basic Information

Catalog Number:

10745-1-AP

Size:

150ul, Concentration: 900 ug/ml by Nanodrop;

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG1199

GenBank Accession Number:

BC011603

GeneID (NCBI):

5970

UNIPROT ID:

Q04206

Full Name:

v-rel reticuloendotheliosis viral oncogene homolog A (avian)

Calculated MW:

65 kDa

Observed MW:

65 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:1000-1:6000

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

IHC 1:50-1:500

IF/ICC 1:50-1:500

Applications

Tested Applications:

WB, IHC, IF/ICC, IP, ELISA

Cited Applications:

WB, IHC, IF, IP, CoIP, ChIP, ELISA

Species Specificity:

human, mouse, rat

Cited Species:

human, canine, monkey, chicken, bovine, hamster, goat, fish, bombyx mori, duck

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: A431 cells, Jurkat cells, MCF-7 cells, K-562 cells, Hek-293 cells, HeLa cells, Raji cells, NIH/3T3 cells

IP: HeLa cells,

IHC: human breast cancer tissue, human colon tissue, human liver cancer tissue, human stomach tissue, mouse colon tissue

IF/ICC: HepG2 cells, TNF alpha treated HT-1080 cells

Background Information

Nuclear factor κ B (NF-κB) is a sequence-specific DNA-binding protein complex which regulates the expression of viral genomes, including the human immunodeficiency virus, and a variety of cellular genes, particularly those involved in immune and inflammatory responses. The members of the NF-κB family in mammalian cells include the proto-oncogene c-Rel, p50/p105 (NFκB1), p65 (RelA), p52/p100 (NFκB2), and RelB. All of these proteins share a conserved 300-amino acid region known as the Rel homology domain which is responsible for DNA binding, dimerization, and nuclear translocation of NF-κB. The p65 subunit is a major component of NF-κB complexes and is responsible for trans-activation. NF-κB heterodimeric p65-p50 and p65-c-Rel complexes are transcriptional activators. The NF-κB p65-p65 complex appears to be involved in invasion-mediated activation of IL-8 expression. The inhibitory effect of IκB upon NF-κB the cytoplasm is exerted primarily through the interaction with p65. p65 shows a weak DNA-binding site which could contribute directly to DNA binding in the NF-κB complex. It associates with chromatin at the NF-κB promoter region via association with DDX1. This antibody is a rabbit polyclonal antibody raised against residues near the N terminus of human RELA.

Notable Publications

Author	Pubmed ID	Journal	Application
Ji Xing	36230734	Cancers (Basel)	WB
Chang Liu	36230117	Foods	WB
Yanliang Wu	34601083	J Ethnopharmacol	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




WB result of RELA, p65 antibody (10745-1-AP, 1:500) with si-Control and si-RELA,p65 transfected HeLa cells.




Various lysates were subjected to SDS PAGE followed by western blot with 10745-1-AP (NF-kB p65 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 10745-1-AP (NF-kB p65 antibody) at dilution of 1:200 (under 0x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using 10745-1-AP (p65; RELA antibody) at dilution of 1:100 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).




IP result of anti-NF-kB p65 (IP:10745-1-AP, 4ug; Detection:10745-1-AP 1:6000) with HeLa cells lysate 1280 ug.



Immunohistochemical analysis of paraffin-embedded mouse colon tissue slide using 10745-1-AP (NF-kB p65 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human colon tissue slide using 10745-1-AP (NF-kB p65 antibody) at dilution of 1:500 (under 20x lens). Heat mediated antigen retrieval with Sodium Citrate buffer (pH 6.0).



Immunofluorescent analysis of (4% PFA) fixed TNF alpha treated HT-1080 cells using NF-kB p65 antibody (10745-1-AP) at dilution of 1:400 and Multi-rAb CoraLite® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002), CL594-phalloidin (red).