For Research Use Only

RHOA Polyclonal antibody

Catalog Number: 10749-1-AP

Featured Product

154 Publications



Basic Information

Catalog Number: GenBank Accession Number:

10749-1-AP BC005976 GeneID (NCBI): Size:

150ul , Concentration: 600 ug/ml by

Nanodrop: **UNIPROT ID:** P61586 Rabbit

Isotype: ras homolog gene family, member A

IgG Calculated MW:

Immunogen Catalog Number: 22 kDa

AG1141 Observed MW:

22 kDa

Full Name:

Applications

Tested Applications:

WB, IF/ICC, IP, ELISA Cited Applications: WB, IHC, IF, IP, CoIP Species Specificity:

human, mouse, rat **Cited Species:**

human, mouse, rat, pig, canine

Positive Controls:

WB: HeLa cells, MCF-7 cells, rat brain tissue, PC-12 cells, HL-60 cells, mouse brain tissue, HUVEC cells,

Purification Method:

WB 1:1000-1:8000

IF/ICC 1:200-1:800

protein lysate

Antigen affinity purification

IP 0.5-4.0 ug for 1.0-3.0 mg of total

Recommended Dilutions:

NIH/3T3 cells, fetal human brain tissue

IP: mouse brain tissue, IF/ICC: HeLa cells, K-562 cells

Background Information

RhoA is a member of the Rho family of small GTPases, which cycle between inactive GDP-bound and active GTPbound states and function as molecular switches in signal transduction cascades. Rho proteins promote reorganization of the actin cytoskeleton and regulate cell shape, attachment, and motility. Overexpression of RhoA is associated with tumor cell proliferation and metastasis. RhoA signalling is critical to many cellular processes including migration, mechanotransduction, and is often disrupted in carcinogenesis.

Notable Publications

Author	Pubmed ID	Journal	Application
Lichan Mao	36247873	Anal Cell Pathol (Amst)	WB
Kai Cui	27678468	Asian J Androl	WB,IHC
Jing Wang	26409448	Tumour Biol	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 10749-1-AP (RHOA antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




Various lysates were subjected to SDS PAGE followed by western blot with 10749-1-AP (RHOA antibody) at dilution of 1:4000 incubated at room temporature for 1.5 hours.



IP result of anti-RHOA (IP:10749-1-AP, 4ug; Detection:10749-1-AP 1:4000) with mouse brain tissue lysate 2100 ug.



Immunofluorescent analysis of (4% PFA) fixed K-562 cells using RHOA antibody (10749-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2).



Immunofluorescent analysis of (-20°C Methanol) fixed HeLa cells using RHOA antibody (10749-1-AP) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (5A00013-2).