

For Research Use Only

UCP3 Polyclonal antibody

Catalog Number: 10750-1-AP

15 Publications



Basic Information

Catalog Number: 10750-1-AP	GenBank Accession Number: BC008392	Purification Method: Antigen affinity purification
Size: 150ul, Concentration: 750 µg/ml by Nanodrop and 400 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 7352	Recommended Dilutions: WB 1:500-1:2000 IHC 1:50-1:500
Source: Rabbit	Full Name: uncoupling protein 3 (mitochondrial, proton carrier)	
Isotype: IgG	Calculated MW: 34 kDa	
Immunogen Catalog Number: AG1134	Observed MW: 30-34 kDa	

Applications

Tested Applications:

IHC, WB, ELISA

Cited Applications:

IHC, WB

Species Specificity:

human, mouse, rat

Cited Species:

human, rat, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: mouse skeletal muscle tissue, SGC-7901 cells, rat skeletal muscle tissue

IHC: human skeletal muscle tissue, human colon cancer tissue

Background Information

Notable Publications

Author	Pubmed ID	Journal	Application
María Rodríguez-Hidalgo	30228311	Sci Rep	WB
Zimengwei Ye	36160420	Front Pharmacol	WB
Sijie Wang	36216077	Physiol Behav	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 10750-1-AP (UCP3 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



mouse skeletal muscle tissue were subjected to SDS PAGE followed by western blot with 10750-1-AP (UCP3 Antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human skeletal muscle tissue slide using 10750-1-AP (UCP3 Antibody) at dilution of 1:200 (under 10x lens).



Immunohistochemical analysis of paraffin-embedded human skeletal muscle tissue slide using 10750-1-AP (UCP3 Antibody) at dilution of 1:200 (under 40x lens).