### For Research Use Only

# **UQCRB** Polyclonal antibody

Catalog Number: 10756-1-AP

**Featured Product** 

**14 Publications** 



**Basic Information** 

Catalog Number: 10756-1-AP

GenBank Accession Number:

**Purification Method:** Antigen affinity purification

Size:

BC005230 GeneID (NCBI):

Recommended Dilutions:

150ul, Concentration: 500 µg/ml by

7381

WB 1:500-1:2000

Nanodrop and 300 µg/ml by Bradford Full Name: method using BSA as the standard;

ubiquinol-cytochrome c reductase

IP 0.5-4.0 ug for IP and 1:500-1:1000

binding protein Calculated MW:

Observed MW:

14 kDa

for WB IHC 1:50-1:500

Rabbit Isotype: IgG

Immunogen Catalog Number:

AG1127

14 kDa

**Applications** 

**Tested Applications:** 

IHC, IP, WB, ELISA

**Cited Applications:** 

IHC, WB

Species Specificity: human, mouse, rat

**Cited Species:** 

human, mouse

**Positive Controls:** 

WB: mouse heart tissue, rat heart tissue

IP: mouse heart tissue.

IHC: human breast cancer tissue, human kidney tissue,

human ovary tumor tissue

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

## **Background Information**

UQCRB(ubiquinol-cytochrome c reductase binding protein) is also named as UQBP, complex III subunit VII and belongs to the UQCRB/QCR7 family. It is a 13.4 kDa subunit of mitochondrial complex III, plays a crucial role in hypoxia-induced angiogenesis via mitochondrial reactive oxygen species (ROS)-mediated signaling(PMID:21215626). This protein has 2 isoforms produced by alternative splicing.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Xiaowen Xu	36071491	Proteome Sci	WB
Yingyan Han	27239088	Dis Markers	WB
Qing Zhang	35594990	Free Radic Biol Med	WB

Storage


Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


## **Selected Validation Data**



Various lysates were subjected to SDS PAGE followed by western blot with 10756-1-AP (UQCRB antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 10756-1-AP (UQCRB antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP Result of anti-UQCRB (IP:10756-1-AP, 3ug; Detection:10756-1-AP 1:600) with mouse heart tissue lysate 3200ug.