

For Research Use Only

COG1 Polyclonal antibody

Catalog Number: 10767-1-AP



Basic Information

Catalog Number: 10767-1-AP	GenBank Accession Number: BC021985	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 550 µg/ml by Nanodrop;	GeneID (NCBI): 9382	Recommended Dilutions: WB 1:1000-1:8000 IHC 1:200-1:800
Source: Rabbit	Full Name: component of oligomeric golgi complex 1	
Isotype: IgG	Calculated MW: 109 kDa	
Immunogen Catalog Number: AG1203	Observed MW: 109 kDa	

Applications

Tested Applications: IHC, WB, ELISA	Positive Controls:
Species Specificity: human	WB : U-251 cells, PC-3 cells, U2OS cells IHC : human ovary tumor tissue,

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

COG1 is one protein of the COG complex which is a heterooctameric complex that regulates intraGolgi trafficking and the integrity of the Golgi compartment in eukaryotic cells (PMID: 16537452). It is reported that this protein is required for steps in the normal medial and trans-Golgi-associated processing of glycoconjugates and plays a role in the organization of the Golgi-localized complex. COG1 has a calculated molecular weight of 109 kDa.

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 10767-1-AP (COG1 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human ovary tumor tissue slide using 10767-1-AP (COG1 antibody) at dilution of 1:400 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human ovary tumor tissue slide using 10767-1-AP (COG1 antibody) at dilution of 1:400 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Various lysates were subjected to SDS PAGE followed by western blot with 10767-1-AP (COG1 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.