

For Research Use Only

# Adrenomedullin Polyclonal antibody



Catalog Number: 10778-1-AP

4 Publications

## Basic Information

<b>Catalog Number:</b> 10778-1-AP	<b>GenBank Accession Number:</b> BC015961	<b>Purification Method:</b> Antigen affinity purification
<b>Size:</b> 150ul , Concentration: 600 µg/ml by Nanodrop and 353 µg/ml by Bradford method using BSA as the standard;	<b>GeneID (NCBI):</b> 133	<b>Recommended Dilutions:</b> WB 1:500-1:2000 IHC 1:50-1:500 IF 1:10-1:100
<b>Source:</b> Rabbit	<b>UNIPROT ID:</b> P35318	
<b>Isotype:</b> IgG	<b>Full Name:</b> adrenomedullin	
<b>Immunogen Catalog Number:</b> AG1197	<b>Calculated MW:</b> 20 kDa	
	<b>Observed MW:</b> 6 kDa	

## Applications

### Tested Applications:

IF, IHC, WB, ELISA

### Cited Applications:

IF, IHC, WB

### Species Specificity:

human, mouse

### Cited Species:

human

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

### Positive Controls:

**WB:** human kidney tissue, fetal human brain tissue, Raji cells, A549 cells, human placenta tissue

**IHC:** human placenta tissue, human pancreas cancer tissue, human kidney tissue, mouse kidney tissue

**IF:** A549 cells,

## Background Information

Adrenomedullin (AM) and proadrenomedullin N-terminal 20 peptide (PAMP) are two small active hormones derived from the expression of a single gene (Adm) that is expressed throughout the GI tract, including the mucosal epithelium, glandular duct cells, neuroendocrine cells, and smooth muscle cells of the GI tract, between the oral cavity and the rectum (PMID:10782362, PMID:27345325). These two peptides coexist in GI cells, where they regulate many physiological functions including vasodilation, angiogenesis, anti-inflammation, organ protection, and tissue repair. AM suppresses inflammatory cytokine production in the intestinal mucosa, improves vascular and lymphatic function, mucosal epithelial repair, and intestinal barrier function in animal models with intestinal inflammation (PMID:27965594, PMID:29311984). Molecular mass species of 18, 14, and 6 kDa were identified in tumor cell lysates and presumably represent AM precursor, processed intermediates, and the authentic peptide, respectively. There is also a 22-kDa immunoreactive species in two cancer cell lines, H720 and MCF-7 (PMID: 8798536).

## Notable Publications

Author	Pubmed ID	Journal	Application
Zhenwei Song	35805068	Cells	IF
Giulia Antoniali	35876890	Cell Mol Life Sci	IHC
Takeshi Sasaki	36479717	Prostate	WB

## Storage

### Storage:

Store at -20°C. Stable for one year after shipment.

### Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com  
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data




Immunohistochemical analysis of paraffin-embedded human placenta tissue slide using 10778-1-AP (Adrenomedullin antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




human kidney tissue were subjected to SDS PAGE followed by western blot with 10778-1-AP (Adrenomedullin antibody at dilution of 1:1000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human placenta tissue slide using 10778-1-AP (Adrenomedullin antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse kidney tissue slide using 10778-1-AP (Adrenomedullin antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded mouse kidney tissue slide using 10778-1-AP (Adrenomedullin antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




A549 cells were subjected to SDS PAGE followed by western blot with 10778-1-AP (Adrenomedullin antibody at dilution of 1:1000 incubated at room temperature for 1.5 hours.




Raji cells were subjected to SDS PAGE followed by western blot with 10778-1-AP (Adrenomedullin antibody at dilution of 1:1000 incubated at room temperature for 1.5 hours.



human placenta tissue were subjected to SDS PAGE followed by western blot with 10778-1-AP (Adrenomedullin antibody at dilution of 1:300 incubated at room temperature for 1.5 hours.



fetal human brain tissue were subjected to SDS PAGE followed by western blot with 10778-1-AP (Adrenomedullin antibody at dilution of 1:300 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Ethanol)  
fixed A549 cells using 10778-1-AP  
(Adrenomedullin antibody) at dilution of 1:25 and  
Alexa Fluor 488-conjugated AffiniPure Goat Anti-  
Rabbit IgG(H+L).