For Research Use Only

TDP-43 Polyclonal antibody

Catalog Number: 10782-2-AP

Featured Product

1434 Publications



Basic Information

Catalog Number: GenBank Accession Number:

10782-2-AP BC001487
Size: Genel D (NCBI):
150ul , Concentration: 600 μg/ml by 23435

Nanodrop; Full Name:
Source: TAR DNA binding protein

Rabbit Calculated MW:
Isotype: 43 kDa
IgG Observed MW:
44 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:2000-1:8000 IF 1:3000-1:12000

WB 1:2000-1:10000

Applications

Tested Applications:

FC, IF, IHC, IP, WB, ELISA

Cited Applications:

ChIP, CoIP, ELISA, IEM, IF, IHC, IP, RIP, WB

Species Specificity:

human, mouse, rat, zebrafish

Cited Species:

human, chicken, horse, Yeast, rat, seal, mouse,

monkey, zebrafish, hamster

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: SH-SY5Y cells, HeLa cells, C2C12 cells, Neuro-2a

cells

IP: HeLa cells,

IHC: human gliomas tissue, human brain (FTLD-U)

tissue, mouse brain tissue

IF: HeLa cells,

Background Information

The TARDBP gene encodes the TDP-43 protein, initially found to repress HIV-1 transcription by binding TAR DNA. TDP-43 has since been shown to bind RNA as well as DNA, and have multiple functions in transcriptional repression, translational regulation and pre-mRNA splicing. For instance, it is reported to regulate alternate splicing of the CTFR gene. In 2006 Neumann et al. found that hyperphosphorylated, ubiquitinated and/or cleaved forms of TDP-43, collectively known as pathological TDP-43, play a major role in the disease mechanisms of ubiquitin-positive, tau-and alpha-synuclein-negative frontotemporal dementia (FTLD-U) and in amyotrophic lateral sclerosis (ALS). Proteintech's 10782-2-AP antibody is a rabbit polyclonal antibody recognizing N-terminal TDP-43. It recognizes the intact 43 kDa protein as well as all posttranslationally modified and truncated forms in multiple applications. Various forms of TDP-43 exist, including 18-35 kDa of cleaved C-terminal fragments, 45-50 kDa phospho-protein, 55 kDa glycosylated form, 75 kDa hyperphosphorylated form, and 90-300 kDa cross-linked form. (17023659, 19823856, 21666678, 22193176) Recently TDP-43 has been reported to be overexpressed in triple negative breast cancer (TNBC) and it may be a potential target for TNBC diagnosis and drug design. (29581274)

Notable Publications

Author	Pubmed ID	Journal	Application
Kasey L Jackson	26445725	Mol Ther Methods Clin Dev	WB
Jinwal Umesh K UK	19793966	J Neurosci	WB
Rudolf Hergesheimer	33003404	Cells	

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


*** 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:


T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com W: ptglab.com This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



40X of FTLD-U case stained by 10782-2-AP and 60019-2-Ig, showing dystrophic neurites. (Figs were provided by Linda K. Kwong).




Various lysates were subjected to SDS PAGE followed by western blot with 10782-2-AP (TDP-43 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human gliomas tissue slide using 10782-2-AP (TDP-43 antibody) at dilution of 1:4000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded human gliomas tissue slide using 10782-2-AP (TDP-43 antibody) at dilution of 1:4000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using TDP-43 antibody (10782-2-AP) at dilution of 1:6000 and Coralite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-phalloidin (red).




Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 10782-2-AP (TDP-43 antibody) at dilution of 1:2000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




WB result of TDP-43 antibody (10782-2-AP; 1:30000; incubated at room temperature for 1.5 hours) with sh-Control and sh-TDP-43 transfected HeLa cells.



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 10782-2-AP (TDP-43 antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-TDP-43(IP:10782-2-AP, 4ug; Detection:10782-2-AP 1:50000) with HeLa cells lysate 1520 ug.



1X10^6 HeLa cells were intracellularly stained with 0.4 ug Anti-Human TDP-43 (10782-2-AP) and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Isotype Control. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).