For Research Use Only

RAB3C Polyclonal antibody

Catalog Number: 10788-1-AP

5 Publications



Basic Information

Catalog Number: GenBank Accession Number:

10788-1-AP Antigen affinity purification BC013033 GeneID (NCBI): Recommended Dilutions: WB 1:2000-1:10000

150ul, Concentration: 550 µg/ml by 115827 Nanodrop;

RAB3C, member RAS oncogene family for WB Source:

Rabbit Calculated MW: Isotype: 26 kDa IgG

Observed MW: Immunogen Catalog Number: 28 kDa

Positive Controls:

WB: mouse brain tissue, human brain tissue, mouse

Purification Method:

IHC 1:50-1:500

IP 0.5-4.0 ug for IP and 1:200-1:1000

lung tissue, rat brain tissues IP: mouse brain tissue.

IHC: mouse brain tissue,

Applications

Tested Applications: IHC, IP, WB, ELISA

Cited Applications: IF, IHC, WB

Species Specificity: human, mouse, rat **Cited Species:**

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

human, mouse

Background Information

Rab proteins are small molecular weight GTPases that control vesicular traffic in eukaryotic cells. A subset of Rab proteins, the Rab3 proteins (Rab3A, Rab3B, Rab3C, and Rab3D) are thought to play an important role in regulated $exocytosis\ of\ vesicles.\ Although\ Rab3\ proteins\ are\ highly\ homologous,\ displaying\ 77-85\%\ amino\ acid\ identity\ with$ greatest variance evident within the N- and C-terminal regions, their subcellular targets and functional roles remain somewhat distinct between them. RT-PCR analysis indicated that human Rab3C was expressed in the human brain, placenta, and lung.

Notable Publications

Author	Pubmed ID	Journal	Application
Maha El Shahawy	28715412	PLoS Genet	IF
Не Ма	36583008	Front Endocrinol (Lausanne)	WB
Wanxue Xu	33311475	Signal Transduct Target Ther	IHC

Storage

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 10788-1-AP (RAB3C antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



IP Result of anti-RAB3C (IP:10788-1-AP, 4ug; Detection:10788-1-AP 1:300) with mouse brain tissue lysate 4000ug.



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 10788-1-AP (RAB3C antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 10788-1-AP (RAB3C antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).