For Research Use Only

Sestrin 2 Polyclonal antibody

Catalog Number:10795-1-AP

Featured Product 209 Publications




Basic Information	Catalog Number: 10795-1-AP	GenBank Accession I BC013304	Number:	Purification Method: Antigen affinity purification				
	Size:	GenelD (NCBI): 83667		Recommended Dilutions: WB 1:2000-1:12000 IP 0.5-4.0 ug for 1.0-3.0 mg of total				
	150ul , Concentration: 750 ug/ml by Nanodrop; Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG1247							
		UNIPROT ID: P58004		IF 0.5-4.0 g foi 1.0-5.0 mg of totat protein lysate IHC 1:200-1:800 IF-P 1:50-1:500 IF/ICC 1:50-1:500				
		Full Name:						
		sestrin 2						
		Calculated MW: 54 kDa						
		Observed MW:						
		54-60 kDa						
Applications	Tested Applications:			trols:				
	WB, IHC, IF/ICC, IF-P, IP, ELISA Cited Applications: WB, IHC, IF, IP, CoIP Species Specificity:		WB : HEK-293 cells, human liver tissue, human brain tissue, NIH/3T3 cells, rat liver tissue, HeLa cells, K-56 cells, mouse kidney tissue, A2780 cells, RAW 264.7 cells					
					human, mouse		IP : K-562 cells,	
					Cited Species: human, mouse, rat, pig, bovine		IHC : human breast cancer tissue, human ovary tumor tissue, human colon cancer tissue	
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0		IF-P : mouse kidney tissue,					
			IF/ICC : HeLa cells,					
		, , ,	vith citrate	IF/ICC : HeLa	cells,			
	Background Information	buffer pH 6.0 SESN is a family of highly conserved oncogenic signaling. SESN2 ,which b inhibit angiogenesis and promote at	, stress-inducible gene elongs to the SESN fan itophagy, underscoring of metabolism and su	es that can defer nily, has been ir g the importance rvival.(PMID:22]	nd the cell against oxidative damage and nplicated as a tumor suppressor that can e of elucidating the molecular mechanisr 363791). Sesn2 encodes a 480 amino acio			
	buffer pH 6.0 SESN is a family of highly conserved oncogenic signaling. SESN2, which b inhibit angiogenesis and promote at by which SESN2 regulates pathways protein which can run as a 60 kDa ba	, stress-inducible gene elongs to the SESN fan itophagy, underscoring of metabolism and su	es that can defer nily, has been ir g the importance rvival.(PMID:22] nide gel electro	nd the cell against oxidative damage and nplicated as a tumor suppressor that can e of elucidating the molecular mechanisr 363791). Sesn2 encodes a 480 amino acio				
	buffer pH 6.0 SESN is a family of highly conserved oncogenic signaling. SESN2, which be inhibit angiogenesis and promote au by which SESN2 regulates pathways protein which can run as a 60 kDa ba Author Put	, stress-inducible gene elongs to the SESN fan itophagy, underscoring of metabolism and su nd on SDS-polyacrylar	es that can defer nily, has been ir g the importance rvival.(PMID:22] nide gel electro	nd the cell against oxidative damage and nplicated as a tumor suppressor that can e of elucidating the molecular mechanisr 363791). Sesn2 encodes a 480 amino acio phoresis(PMID:20878915).				
Background Information Notable Publications	buffer pH 6.0 SESN is a family of highly conserved oncogenic signaling. SESN2, which binhibit angiogenesis and promote at by which SESN2 regulates pathways protein which can run as a 60 kDa ba Author Put Min Peng	, stress-inducible gene elongs to the SESN fan itophagy, underscoring of metabolism and su nd on SDS-polyacrylar omed ID Jourr	es that can defer nily, has been ir g the importance rvival.(PMID:22: nide gel electro nal	nd the cell against oxidative damage and nplicated as a tumor suppressor that can e of elucidating the molecular mechanisr 363791). Sesn2 encodes a 480 amino acid phoresis(PMID:20878915). Application				
	buffer pH 6.0 SESN is a family of highly conserved oncogenic signaling. SESN2, which binhibit angiogenesis and promote at by which SESN2 regulates pathways protein which can run as a 60 kDa ba Author Put Min Peng Scot R Kimball 264	, stress-inducible gene elongs to the SESN fan itophagy, underscoring of metabolism and su nd on SDS-polyacrylar omed ID Journ 259925 Cell	es that can defer nily, has been ir g the importance rvival.(PMID:22: nide gel electro nal	nd the cell against oxidative damage and nplicated as a tumor suppressor that can e of elucidating the molecular mechanisr 363791). Sesn2 encodes a 480 amino acid phoresis(PMID:20878915). Application WB				
	buffer pH 6.0 SESN is a family of highly conserved oncogenic signaling. SESN2, which bi inhibit angiogenesis and promote at by which SESN2 regulates pathways protein which can run as a 60 kDa ba Author Put Min Peng Scot R Kimball 264	l, stress-inducible gene elongs to the SESN fan itophagy, underscoring of metabolism and su nd on SDS-polyacrylar omed ID Jourr 259925 Cell 200964 J Nut 290574 Antic ter shipment.	es that can defer nily, has been ir g the importance rvival.(PMID:22; nide gel electro nal	nd the cell against oxidative damage and nplicated as a tumor suppressor that can e of elucidating the molecular mechanisr 363791). Sesn2 encodes a 480 amino acid phoresis(PMID:20878915). Application WB				


For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 10795-1-AP (Sestrin 2 antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours.




WB result of Sestrin antibody (10795-1-AP, 1:2000) with si-Control and si-Sestrin transfected HEK293 cells.




Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 10795-1-AP (Sestrin 2 antibody) at dilution of 1:400 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed mouse kidney tissue using Sestrin 2 antibody (10795-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunofluorescent analysis of (-20°C Ethanol) fixed HeLa cells using Sestrin 2 antibody (10795-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2).



IP result of anti-Sestrin 2 (IP:10795-1-AP, 3ug; Detection:10795-1-AP 1:1000) with K-562 cells lysate 4000ug.